

IOWA STATE UNIVERSITY

Digital Repository

Graduate Theses and Dissertations

Iowa State University Capstones, Theses and
Dissertations

2013

Obesity and adipose tissue expansion in Ossabaw swine

Richard Joel Faris
Iowa State University

Follow this and additional works at: <https://lib.dr.iastate.edu/etd>

 Part of the [Allergy and Immunology Commons](#), [Human and Clinical Nutrition Commons](#), [Immunology and Infectious Disease Commons](#), [Medical Immunology Commons](#), and the [Physiology Commons](#)

Recommended Citation

Faris, Richard Joel, "Obesity and adipose tissue expansion in Ossabaw swine" (2013). *Graduate Theses and Dissertations*. 12990.
<https://lib.dr.iastate.edu/etd/12990>

This Dissertation is brought to you for free and open access by the Iowa State University Capstones, Theses and Dissertations at Iowa State University Digital Repository. It has been accepted for inclusion in Graduate Theses and Dissertations by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Obesity and adipose tissue expansion in Ossabaw swine

by

Richard J. Faris

A dissertation submitted to the graduate faculty
in partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY

Major: Nutritional Sciences

Program of Study Committee:
Michael E. Spurlock, Major Professor
Donald C. Beitz
Nicholas K. Gabler
James Hollis
Douglas E. Jones

Iowa State University

Ames, Iowa

2013

Copyright © Richard J. Faris, 2013. All rights reserved.

TABLE OF CONTENTS

ABBREVIATIONS	v
LIST OF FIGURES	vii
LIST OF TABLES	viii
ACKNOWLEDGEMENTS	ix
ABSTRACT	x
CHAPTER 1. GENERAL INTRODUCTION	1
Introduction	1
Objective and Specific Aims	3
References	4
CHAPTER 2. REVIEW OF LITERATURE	7
Obesity Prevalence	7
Obesity, Inflammation, and Insulin Resistance	8
Adipose Tissue Expansion	8
Insulin Signaling	10
Tumor Necrosis Factor α	11
Interleukin-6	12
C-Reactive Protein	13
Monocyte Chemoattractant Protein 1	14
Interleukin-8	15
Adiponectin	16
Oxidative Stress	17
Metabolic Endotoxemia	17

Obesity and Adipose Tissue Macrophages	18
Obesity and Fatty Acids	21
Swine Models of Obesity	23
Summary	29
References	30
CHAPTER 3: INFLAMMATION AND RESPONSE TO N-3 FATTY ACIDS IN A PORCINE OBESITY MODEL	62
Abstract	62
Introduction	63
Methods	64
Results	69
Discussion	72
Acknowledgements	76
References	77
CHAPTER 4: ADIPOGENIC POTENTIAL OF OSSABAW ADIPOCYTES PROTECTS AGAINST IL-8 SECRETION	88
Abstract	88
Introduction	89
Methods	91
Results	95
Discussion	97
References	101

CHAPTER 5: ADIPOSE TISSUE DEPOT DIFFERENCES IN MICROARRAY EXPRESSION IN OSSABAW SWINE	112
Abstract	112
Introduction	113
Methods	114
Results	116
Discussion	118
References	120
CHAPTER 6: GENERAL CONCLUSIONS	133
References	137

ABBREVIATIONS

2-NBDG	2-deoxy-[(7-nitro-2,1,3-benzoxadiazol-4-yl)amino]-D-glucose
AdipoR	adiponectin receptor
AMPK	AMP activated protein kinase
BMI	body mass index
BrDU	bromodeoxyuridine
BSA	bovine serum albumin
CD	cluster of differentiation (i.e. CD14)
CCL	chemokine (C-C motif) ligand (i.e. CCL2)
CCR	C-C chemokine receptor (i.e. CCR2)
COX	cyclooxygenase
CXCL	chemokine (C-X-X motif) ligand (i.e. CXCL10)
C/EBP	CCAAT/enhancer binding protein
CRP	C-reactive protein
DGAT	diacylglycerol acyltransferase
DHA	docosahexaenoic acid
EPA	eicosapentaenoic acid
ERK	extracellular signal-related protein kinase
ERp44	endoplasmic reticulum protein 44
FDR	false discovery rate
FTO	fat mass and obesity associated
G3PDH	glycerol-3-phosphate dehydrogenase
GLUT	glucose transporter
HBSS	Hank's buffered salt solution
HOMA-IR	Homeostasis Model of Assessment - Insulin Resistance
IDO	indoleamine 2,3-dioxygenase
IKK β	inhibitor of nuclear factor κ B subunit β
IGF	insulin-like growth factor
IGFBP	insulin-like growth factor binding protein
IL	interleukin
iNOS	inducible nitric oxide synthase
IRS	insulin receptor substrate
Jak	Janus kinase
JNK	Jun N-terminal kinase
LPS	lipopolysaccharide
M1	classically activated macrophages
M2	alternatively activated macrophages
MAPK	mitogen-activated protein kinase
MCP	monocyte chemoattractant protein
MEK	mitogen-activated protein kinase kinase (MAPK/ERK kinase)
MHO	metabolically healthy obese
NADPH	Nicotinamide adenine dinucleotide phosphate
NEFA	non-esterified fatty acids
NF κ B	nuclear factor κ B

NO	nitric oxide
PBS	phosphate buffered saline
PI3K	phosphatidylinositol 3-kinase
PPAR	peroxisome proliferator-activated receptor
PKB	protein kinase B
PUFA	polyunsaturated fatty acids
RNS	reactive nitrogen species
ROS	reactive oxygen species
SNP	single nucleotide polymorphism
SOCS	suppressor of cytokine signaling
STAT	signal transducer and activator of transcription
SVC	stromal vascular cell
Tlr	toll-like receptor
TNF	tumor necrosis factor
TNFR	tumor necrosis factor α receptor
TZD	thiazolidinedione

LIST OF FIGURES

Figure 1. <i>n-3 fatty acids attenuate the high palm oil induced decrease in serum adiponectin, but do not alter AMPK.</i>	84
Figure 2. <i>n-3 fatty acids attenuate high palm oil induced insulin resistance.</i>	85
Figure 3. <i>Insulin resistance is not reflected in adipose tissue S₄₇₃ PKB phosphorylation.</i>	86
Figure 4. <i>High fat diets have fewer macrophages, but phenotype is altered by obesity and depot.</i>	87
Figure 5. <i>Comparison of SQ and VIS adipocyte size in lean and obese Ossabaw Swine.</i>	106
Figure 6. <i>Ossabaw adipocytes have increased clonal expansion and G3PDH activity than Yorkshire adipocytes.</i>	107
Figure 7. <i>Yorkshire adipocytes have more lipid filling than Ossabaw adipocytes.</i>	107
Figure 8. <i>Ossabaw adipocytes express more cellular PPARγ1 than Yorkshire adipocytes.</i>	108
Figure 9. <i>Ossabaw adipocytes have decreased insulin sensitivity compared to Yorkshire adipocytes, but do not respond differently to palmitate or LPS treatment.</i>	109
Figure 10. <i>Palmitate and LPS differentially regulate inflammation in Ossabaw and Yorkshire adipocytes.</i>	110
Figure 11. <i>Ossabaw swine do not have obesity-linked increases in IL-8 or myeloperoxidase activity.</i>	111

LIST OF TABLES

Table 1. <i>Swine subcutaneous adipose tissue macrophage phenotype.</i>	61
Table 2. <i>Ingredient composition of the diet (%) as-fed.</i>	81
Table 3. <i>Dietary fatty acid profile (g/100 g total fatty acid).</i>	82
Table 4. <i>Growth and metabolic parameters of Ossabaw swine at termination.</i>	83
Table 5. <i>Metabolic profile of Ossabaw swine.</i>	124
Table 6. <i>Differentially expressed genes by depot ($Q < 0.01$).</i>	125
Table 7. <i>Gene ontology cellular component: extracellular region gene list^a.</i>	129
Table 8. <i>KEGG pathway: arachidonic acid metabolism gene list^a.</i>	131
Table 9. <i>KEGG pathway: chemokine signaling pathway gene list^a.</i>	132

ACKNOWLEDGEMENTS

First and foremost, I would like to thank my major professor, Dr. Michael E. Spurlock. His guidance, support, and understanding throughout my graduate education has been instrumental to my success. I would also like to extend thanks to my committee members Dr. Donald Beitz, Dr. Douglas Jones, Dr. James Hollis, and Dr. Nicholas Gabler. I want to thank my co-workers, both past and present, for their assistance and continued support. I am especially grateful to Dr. Jennifer Walker-Daniels, Dr. Rebecca Boddicker, and Dr. Douglas Braucher. Without their support, I would not be the scientist I am today. Thank you to Dr. Nicholas Gabler, Dr. John Patience, and Dr. Michael Persia for the use of their laboratory after the Spurlock laboratory fire in August of 2010. Thank you to Martha Jeffrey, Julie Roberts, and Neva Nachtrieb for your assistance, support, and friendship. Finally, thank you to all of my friends and family who have supported me throughout graduate school. Without all of you, I would not be where I am today.

ABSTRACT

Obesity is characterized as a state of chronic low-grade inflammation, which originates in adipose tissue from adipocyte dysfunction as a consequence of adipocyte hypertrophy. Although rodent models have been pivotal to understanding the etiology of obesity and its co-morbidities, and the roles of specific genes, distinct differences between rodents and humans underscore the need for alternative comparative models. Swine are an attractive comparative model because of similar physiology and anatomy with humans. Ossabaw swine are of particular interest due to their “thrifty” genotype and predisposition to multiple risk factors for the metabolic syndrome.

Saturated fatty acids have the distinct ability to induce inflammation and insulin resistance via activation of toll-like receptor 4, while n-3 fatty acids generally reduce inflammation and restore insulin sensitivity. Therefore, we first used Ossabaw swine to investigate the effects of high saturated fat and high saturated fat plus n-3 fatty acids on adipose tissue macrophages, inflammation, and insulin resistance. In this study, despite severe obesity, Ossabaw swine did not develop systemic inflammation. Consumption of the high palm oil diet did decrease serum adiponectin, and this decrease was attenuated by the addition of n-3 fatty acids to the diet. Additionally, we determined that the majority of adipose tissue macrophages are of an anti-inflammatory phenotype. High dietary palm oil did increase the percentage of inflammatory macrophages, and this increase was attenuated by the addition of n-3 fatty acids. Finally, the consumption of high palm oil induced insulin resistance, which was attenuated by the addition of n-3 fatty acids. Thus, in Ossabaw swine, morbid obesity induces insulin resistance, but insulin resistance is not linked to inflammation.

Obesity in humans and rodents causes an increase in subcutaneous adipocyte size of ~2 to 3 fold. Ossabaw subcutaneous adipocyte size was increased ~4.3 fold due to the consumption of a diet high in fat. Because of marked adipocyte hypertrophy coupled with the lack of inflammation in obese Ossabaw adipocytes, we sought to determine if Ossabaw adipocytes were protected against saturated fatty acid-induced inflammation. Therefore, we investigated the differences between Ossabaw and Yorkshire stromal vascular cell-derived adipocytes with respect to differentiation and palmitate-induced inflammation and insulin resistance. In this study, we determined that Ossabaw adipocytes express more peroxisome proliferator-activated receptor γ , have more glycerol-3-phosphate dehydrogenase activity, and higher rates of clonal expansion than Yorkshire adipocytes. These findings indicate that Ossabaw adipocytes differentiate to a greater extent than Yorkshire adipocytes. Palmitate decreased insulin-stimulated glucose uptake in both adipocyte genotypes. However, Ossabaw adipocytes had decreased basal and insulin-stimulated glucose uptake compared with Yorkshire adipocytes, although the fold increase was not different with insulin stimulation. With respect to inflammation, palmitate induced nuclear factor κ B activation equally in both adipocyte genotypes, but only Yorkshire adipocytes secreted interleukin 8 in response to palmitate. Consistent with this, there were no differences in serum or adipose tissue IL-8 concentrations between lean and obese Ossabaw swine. Interleukin 8 induces neutrophil chemotaxis into tissues. Neutrophil infiltration into adipose tissue may be important in the early stages of obesity. These data suggest that differences in Ossabaw adipocytes compared with Yorkshire adipocytes may be instrumental in protecting against inflammation during high fat diet-induced obesity.

Finally, we evaluated global gene expression in subcutaneous and visceral adipose tissue of restricted and *ad libitum* fed Ossabaw swine by using an Affymetrix microarray. Microarray analysis revealed no differentially expressed genes between lean and obese Ossabaw swine. This result coincides with no metabolic alterations induced by obesity relating to glucose homeostasis or serum lipids. There were a total of 1494 genes differentially expressed by adipose tissue depot, of which 777 were annotated. DAVID was used to identify functional categories that contained over-represented genes that were differentially expressed by depot. Extracellular space was identified as a gene ontology cellular component category. Two KEGG pathways were identified by DAVID. These were arachidonic acid metabolism and chemokine signaling pathway.

Collectively, we have demonstrated that dietary factors influence the development of metabolic alterations in Ossabaw swine as evidenced by the induction of insulin resistance by high dietary palm oil, the attenuation of insulin resistance by the addition of n-3 fatty acids, and the lack of differentially expressed genes in the adipose tissue of obese Ossabaw swine when consuming a diet without added fat. Additionally, even in the presence of insulin resistance, there was minimal evidence of inflammation beyond the slight increase in the inflammatory CD16⁺CD14⁺ macrophages in the adipose tissue of Ossabaw swine fed a high dietary palm oil. Palmitate-induced interleukin 8 secretion by Yorkshire adipocytes, but not Ossabaw adipocytes, further supports that Ossabaw swine are resistant to obesity-linked inflammation. The ability of Ossabaw swine to expand adipose tissue may afford protection against the induction of inflammation by diet-induced obesity. Further research into the role of adipose expansion and protection against inflammation is warranted.

CHAPTER 1: GENERAL INTRODUCTION

Introduction

In 2007-2008, it was estimated that 33.9% of the population in the United States is obese (BMI \geq 30) (3), compared with 30.5% in 1999-2000 (4). Obesity and its co-morbidities have placed a significant economic burden on our health care system (20-21; 26). Obesity is linked with a cluster of co-morbidities, that when present, meet the defining criteria for the metabolic syndrome. The metabolic syndrome places individuals at a greater risk for the development of type 2 diabetes (5; 14) and cardiovascular disease (6; 14). Additionally, visceral adiposity and fatty liver associated with obesity pose a greater risk for the development of the metabolic syndrome (9). Because of the increase in prevalence of obesity, there has been increased research effort directed at understanding the etiology of the epidemic and identifying potential interventions.

Adipose tissue, once considered just an energy storage tissue, is now recognized as a fully functional endocrine and immune tissue. This tissue produces hormones, cytokines, and chemokines that modulate metabolic and immunological pathways, both centrally and in peripheral tissues. These molecules are termed adipokines. Chronic positive energy balance induces adipose tissue expansion via adipocyte hypertrophy and/or preadipocyte hyperplasia. Adipocyte hypertrophy and dysregulation leads to an altered adipokine profile that precedes macrophage infiltration (8). Adipose remodeling, or turnover of these “enlarged” adipocytes leads to the recruitment of macrophages, which assist in the necrosis of adipocytes (23). Adipocyte hypertrophy and macrophage infiltration alter adipose tissue production and secretion of adipokines such as tumor necrosis factor (TNF) α , interleukin 6 (IL-6), C-reactive protein (CRP), monocyte chemoattractant protein 1 (MCP-1), IL-8, and adiponectin.

Collectively, obesity-induced dysregulation of these adipokines can induce adipose tissue insulin resistance, leading to ectopic lipid deposition, particularly in the liver (2), muscle (18), and pancreas (22).

Consumption of saturated fat can contribute to adipose tissue inflammation. Saturated fatty acids are a ligand for toll-like receptor 4 (11), the innate immune receptor for lipopolysaccharide, and leads to the activation of nuclear factor κ B (NF κ B) and subsequent inflammation. Conversely, n-3 fatty acids are generally considered anti-inflammatory. Eicosapentaenoic acid and docosahexaenoic acid are the n-3 fatty acids with biological activity. These n-3 fatty acids can inhibit NF κ B (10; 17) and activate peroxisome proliferator-activated receptor γ (1).

Nearly a decade after the link between obesity, inflammation, and insulin resistance was established (7), two laboratories independently discovered that obesity is associated with macrophage accumulation in adipose tissue (27-28). Consistent with this, both body mass and adipocyte size are positively correlated with the number of macrophages present in adipose tissue (27).

Mice have been pivotal to the determination of the role of macrophages in adipose tissue. However, distinct differences between human and murine immune function, particularly regarding the role of CRP as an acute phase protein (24) and nitric oxide production in macrophages (15; 19; 25), limit mice as an immunological model for humans. Swine are increasingly used as a biomedical research model, in large part because of their similar physiology and anatomy with humans (12). Recently, swine have been used to study a number of issues relating to obesity, including cardiovascular disease, steatohepatitis, and insulin signaling. Ossabaw swine are of particular interest, given their “thrifty” genotype due

to genetic isolation (13) and ability to develop all of the risk factors associated with the metabolic syndrome (16).

Despite evidence demonstrating the value of swine as a model of obesity, there has not been an evaluation of adipose tissue inflammation in swine under conditions of diet-induced obesity, particularly regarding macrophage infiltration and induction of inflammation via saturated fatty acids, and the inhibition by n-3 fatty acids. Consequently, there is a clear need to *demonstrate adipose tissue inflammation and macrophage infiltration in Ossabaw swine, and to determine the effects of saturated and n-3 fatty acids on inflammation to further evaluate swine as a comparative model for human obesity.*

Objective and Specific Aims

The *objective* of this study was to determine the extent to which Ossabaw swine develop adipose tissue inflammation and macrophage infiltration, and its effects on insulin sensitivity under conditions of diet-induced obesity. The *central hypothesis* was that diet-induced obesity will cause adipose tissue inflammation concomitant with macrophage infiltration in Ossabaw swine. The *rationale* for our work was to establish a link between adipose tissue inflammation and obesity-linked metabolic consequences in an alternative comparative model for human obesity. Our *specific aims* include i) to determine the effect of saturated and n-3 fatty acids on insulin resistance and macrophage phenotype in Ossabaw swine, ii) to determine the effect of saturated fatty acids on Ossabaw and Yorkshire adipocytes in regards to inflammation and insulin resistance, and iii) identify specific gene categories and pathways in Ossabaw swine adipose tissue that are involved in adipose tissue expansion due to diet-induced obesity.

References

1. Banga A, Unal R, Tripathi P, Pokrovskaya I, Owens RJ, Kern PA, Ranganathan G. 2009. Adiponectin translation is increased by the PPARgamma agonists pioglitazone and omega-3 fatty acids. *Am J Physiol Endocrinol Metab* 296:E480-489.
2. Donnelly KL, Smith CI, Schwarzenberg SJ, Jessurun J, Boldt MD, Parks EJ. 2005. Sources of fatty acids stored in liver and secreted via lipoproteins in patients with nonalcoholic fatty liver disease. *J Clin Invest* 115:1343-1351.
3. Flegal KM, Carroll MD, Ogden CL, Curtin LR. 2010. Prevalence and trends in obesity among US adults, 1999-2008. *JAMA* 303:235-241.
4. Flegal KM, Carroll MD, Ogden CL, Johnson CL. 2002. Prevalence and trends in obesity among US adults, 1999-2000. *JAMA* 288:1723-1727.
5. Ford ES, Schulze MB, Pischon T, Bergmann MM, Joost HG, Boeing H. 2008. Metabolic syndrome and risk of incident diabetes: findings from the European Prospective Investigation into Cancer and Nutrition-Potsdam Study. *Cardiovasc Diabetol* 7:35.
6. Galassi A, Reynolds K, He J. 2006. Metabolic syndrome and risk of cardiovascular disease: a meta-analysis. *Am J Med* 119:812-819.
7. Hotamisligil GS, Shargill NS, Spiegelman BM. 1993. Adipose expression of tumor necrosis factor-alpha: direct role in obesity-linked insulin resistance. *Science* 259:87-91.
8. Jiao P, Chen Q, Shah S, Du J, Tao B, Tzamelis I, Yan W, Xu H. 2009. Obesity-related upregulation of monocyte chemotactic factors in adipocytes: involvement of nuclear factor-kappaB and c-Jun NH2-terminal kinase pathways. *Diabetes* 58:104-115.
9. Kim LJ, Nalls MA, Eiriksdottir G, Sigurdsson S, Launer LJ, Koster A, Chaves PH, Jonsdottir B, Garcia M, Gudnason V, Harris TB. 2011. Associations of visceral and liver fat with the metabolic syndrome across the spectrum of obesity: the AGES-Reykjavik study. *Obesity (Silver Spring)* 19:1265-1271.
10. Lee JY, Plakidas A, Lee WH, Heikkinen A, Chanmugam P, Bray G, Hwang DH. 2003. Differential modulation of Toll-like receptors by fatty acids: preferential inhibition by n-3 polyunsaturated fatty acids. *J Lipid Res* 44:479-486.
11. Lee JY, Sohn KH, Rhee SH, Hwang D. 2001. Saturated fatty acids, but not unsaturated fatty acids, induce the expression of cyclooxygenase-2 mediated through Toll-like receptor 4. *J Biol Chem* 276:16683-16689.
12. Litten-Brown JC, Corson AM, Clarke L. 2010. Porcine models for the metabolic syndrome, digestive and bone disorders: a general overview. *Animal* 4:899-920.

13. Mayer JJ, Brisbin IL. 1991. *Wild pigs in the United States : their history, comparative morphology, and current status*. Athens: University of Georgia Press.
14. Meigs JB, Wilson PW, Fox CS, Vasan RS, Nathan DM, Sullivan LM, D'Agostino RB. 2006. Body mass index, metabolic syndrome, and risk of type 2 diabetes or cardiovascular disease. *J Clin Endocrinol Metab* 91:2906-2912.
15. Mestas J, Hughes CC. 2004. Of mice and not men: differences between mouse and human immunology. *J Immunol* 172:2731-2738.
16. Neeb ZP, Edwards JM, Alloosh M, Long X, Mokelke EA, Sturek M. 2010. Metabolic syndrome and coronary artery disease in Ossabaw compared with Yucatan swine. *Comp Med* 60:300-315.
17. Novak TE, Babcock TA, Jho DH, Helton WS, Espat NJ. 2003. NF-kappa B inhibition by omega -3 fatty acids modulates LPS-stimulated macrophage TNF-alpha transcription. *Am J Physiol Lung Cell Mol Physiol* 284:L84-89.
18. Phillips DI, Caddy S, Ilic V, Fielding BA, Frayn KN, Borthwick AC, Taylor R. 1996. Intramuscular triglyceride and muscle insulin sensitivity: evidence for a relationship in nondiabetic subjects. *Metabolism* 45:947-950.
19. Roshick C, Wood H, Caldwell HD, McClarty G. 2006. Comparison of gamma interferon-mediated antichlamydial defense mechanisms in human and mouse cells. *Infect Immun* 74:225-238.
20. Ryan JG. 2009. Cost and policy implications from the increasing prevalence of obesity and diabetes mellitus. *Gend Med* 6 Suppl 1:86-108.
21. Shamseddeen H, Getty JZ, Hamdallah IN, Ali MR. 2011. Epidemiology and economic impact of obesity and type 2 diabetes. *Surg Clin North Am* 91:1163-1172, vii.
22. Shimabukuro M, Zhou YT, Levi M, Unger RH. 1998. Fatty acid-induced beta cell apoptosis: a link between obesity and diabetes. *Proc Natl Acad Sci U S A* 95:2498-2502.
23. Strissel KJ, Stancheva Z, Miyoshi H, Perfield JW, 2nd, DeFuria J, Jick Z, Greenberg AS, Obin MS. 2007. Adipocyte death, adipose tissue remodeling, and obesity complications. *Diabetes* 56:2910-2918.
24. Szalai AJ, McCrory MA. 2002. Varied biologic functions of C-reactive protein: lessons learned from transgenic mice. *Immunol Res* 26:279-287.
25. Thoma-Uszynski S, Stenger S, Takeuchi O, Ochoa MT, Engele M, Sieling PA, Barnes PF, Rollinghoff M, Bolcskei PL, Wagner M, Akira S, Norgard MV, Belisle

- JT, Godowski PJ, Bloom BR, Modlin RL. 2001. Induction of direct antimicrobial activity through mammalian toll-like receptors. *Science* 291:1544-1547.
26. Wang YC, McPherson K, Marsh T, Gortmaker SL, Brown M. 2011. Health and economic burden of the projected obesity trends in the USA and the UK. *Lancet* 378:815-825.
 27. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW, Jr. 2003. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* 112:1796-1808.
 28. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS, Tartaglia LA, Chen H. 2003. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest* 112:1821-1830.

CHAPTER 2: LITERATURE REVIEW

Obesity Prevalence

Obesity is an increasing trend in the United States and other developed countries. In 2007-2008, it was estimated that 33.9% of the population in the United States is obese (BMI ≥ 30) and 68.3% is overweight (BMI ≥ 25) (92). In comparison, the estimates were 30.5% and 64.5%, respectively, in 1999-2000 (93). Obesity and its co-morbidities have placed a significant economic burden on our health care system, and costs are continuing to rise (237; 242; 279). In part, due to public health concern and economic cost, obesity and its co-morbidities have become a large research focus to understand the etiology of the epidemic and to identify potential interventions. However, a myriad of factors, including genotype and environment can contribute to the increasing prevalence of obesity.

Common polymorphisms in the FTO (fat mass and obesity associated) gene and near the melanocortin 4 receptor gene have been associated with the increased risk of obesity. These polymorphisms are associated with increased fat mass and increased risk of obesity (44; 70; 99; 176). Additionally, the presence of multiple risk alleles has an additive effect on the risk of obesity (44). However, the presence of single nucleotide polymorphisms (SNP) cannot completely account for the increased incidence in obesity. For example, the SNP rs1421085 in the FTO gene has an allelic frequency of 0.460 for the risk allele from a US population of European descent, but only a frequency of 0.066 for a United States population of African descent (HapMap). Clearly, more than genetic susceptibility contributes to the increase in obesity prevalence. Thus, genetic susceptibility, in conjunction with urbanization and poor diets has facilitated the increase in obesity prevalence (268).

Obesity is associated with a cluster of co-morbidities that, when present, is defined as the metabolic syndrome. The presence of the metabolic syndrome places individuals at a greater risk for the development of type 2 diabetes (98; 191) and cardiovascular disease (105; 191). The International Diabetes Federation and the American Heart Association/National Heart, Lung, and Blood Institute define the metabolic syndrome as meeting three out of the five following criteria: waist circumference greater than 102 cm for men and 88 cm for women, serum triglycerides greater than 150 mg/dL, HDL cholesterol less than 40 mg/dL for men and 50 mg/dL for women, fasting blood glucose greater than 100 mg/dL, and hypertension with systolic blood pressure greater than 130 mm Hg or diastolic blood pressure greater than 85 mm Hg (7). Additionally, visceral adiposity and fatty liver associated with obesity place individuals at a greater risk for the development of the metabolic syndrome (146).

Obesity, Inflammation, and Insulin Resistance

Adipose Tissue Expansion

Adipose tissue, once considered just an energy storage tissue, is now recognized as a fully functional endocrine and immune organ. This tissue produces a myriad of hormones, cytokines, and chemokines that are capable of modulating metabolic and immunological pathways, both centrally and in peripheral tissues. These molecules, collectively referred to as adipokines, include cytokines, chemokines, adhesion molecules, angiogenic factors, and growth factors that regulate food intake, metabolism, vascularization, reproduction, adipose development, inflammation, and inflammatory cell recruitment (53; 116; 178). Nearly all of the co-morbidities associated with obesity can be linked to chronic inflammation (9; 197; 307).

Chronic positive energy balance induces adipose tissue expansion via adipocyte hypertrophy and/or preadipocyte hyperplasia. Evidence suggests that adipose tissue hypertrophy is genetically limited and varies by depot (69). Adipocyte hypertrophy and dysregulation leads to an altered adipokine profile which precedes macrophage infiltration (131). Death and necrosis of adipocytes, as well as pro-inflammatory adipokine secretion, is positively linked to adipocyte size (54; 129; 250). Adipose remodeling, or turnover of these “enlarged” adipocytes, leads to the recruitment of macrophages, which assist in the necrosis of adipocytes (259). Adipokine-mediated inflammation furthers the recruitment of macrophages and leads to increased inflammation (137; 287). Adipocyte hypertrophy and macrophage infiltration alter adipose tissue production and secretion of adipokines such as tumor necrosis factor (TNF) α , interleukin 6 (IL-6), C-reactive protein (CRP), monocyte chemoattractant protein 1 (MCP-1), IL-8, and adiponectin as well as others.

Adipose tissue inflammation is a potential link between insulin resistance and the eventual development of type 2 diabetes mellitus. It is estimated that 80 % of diabetics are considered overweight and 49 % are obese (204). Thus, there is a strong link between obesity and the development of type 2 diabetes mellitus.

Adipose tissue insulin resistance results in aberrant regulation of the balance between lipogenesis and lipolysis (18). As a consequence, lipolysis is favored and circulating concentrations of free fatty acids are increased. Additionally, postprandial lipid storage is impaired, which further contributes to dyslipidemia (189). As a result, ectopic lipid deposition occurs particularly in the liver (73), muscle (220), and pancreas (246). Ectopic lipid deposition in the liver and muscle is associated with insulin resistance (151; 181), potentially increasing hepatic glucose production and decreasing skeletal muscle glucose

uptake. Furthermore, ectopic lipid deposition in the pancreas can lead to a decrease in β -cell function (246), exacerbating insulin resistance and leading to the development of frank diabetes.

However, some obese individuals do not display the metabolic phenotype often associated with obesity (141; 248). These individuals are termed metabolically healthy but obese (MHO). MHO individuals retain high levels of insulin sensitivity, and have a favorable lipid and inflammatory profile (3; 35; 140; 254). Body composition may be a determining factor in the MHO phenotype. As such, MHO individuals have less visceral (35; 128; 140; 247; 288) and liver (254) fat compared to their metabolically abnormal counterparts. Potentially, the ability to expand adipose tissue may be a fundamental component to the favorable metabolic profile in MHO individuals. In *ob/ob* mice, the induction of adipose tissue expansion attenuates insulin resistance and improves systemic inflammation normally seen in this model, despite morbid obesity (145). This improvement was accompanied by smaller adipocytes and decreased ectopic lipid deposition, suggesting that limits on the ability of adipose tissue to expand cause adipocyte hypertrophy and leads to ectopic lipid deposition that may be causal to the development of insulin resistance in obese individuals.

Insulin Signaling

Insulin is responsible for multiple biological effects, including stimulation of glucose uptake, lipogenesis, protein synthesis, glycogen synthesis, gene transcription, and growth and differentiation. Upon binding of insulin to the insulin receptor, the receptor undergoes auto-phosphorylation of tyrosine residues. Auto-phosphorylation allows the docking and tyrosine phosphorylation of insulin receptor substrate (IRS) proteins (i.e., IRS-1, IRS-2). Tyrosine

phosphorylation of IRS allows the binding of several adaptor proteins including phosphatidylinositol 3-kinase (PI3K). PI3K catalyzes the conversion of phosphatidylinositol 4,5-bisphosphate to phosphatidylinositol 3,4,5-triphosphate, leading to the down-stream activation of protein kinase B (PKB, i.e., Akt) by phosphorylation of threonine 308 and serine 473, and eventual translocation of GLUT4 to the plasma membrane (as reviewed by Taha and Klip) (266).

Tumor Necrosis Factor α

TNF α is an adipokine that can act both locally and systemically (142). In humans, the production of TNF α is elevated in obesity (142-143; 186), but decreases with weight loss (63; 143). TNF α has two receptors, TNFR1 and TNFR2. TNFR1 is the major receptor in adipose tissue, leading to the activation of nuclear factor κ B (NF κ B), and mitogen-activated protein kinases (MAPK) such as extracellular signal-regulated protein kinase (ERK), p38 MAPK, and c-Jun N-terminal kinase (JNK) (45). In obesity, elevated TNF α concentrations are directly linked to insulin resistance (123) and cardiovascular disease (249). Additionally, TNF α up-regulates the expression of pro-inflammatory adipokines (186; 278), and down-regulates the expression of anti-inflammatory adipokines, such as adiponectin (37; 233; 278). At the expense of inflammation and insulin resistance, increased TNF α concentrations during obesity may be viewed as a negative regulator of adiposity and weight gain by increasing lipolysis, inhibiting preadipocyte differentiation, and inducing adipocyte apoptosis (222).

Molecular mechanisms of TNF α -mediated inhibition of insulin signaling involve serine phosphorylation and transcriptional regulation. TNF α inhibits tyrosine

phosphorylation of the insulin receptor and IRS (86), and reduces insulin receptor, IRS-1, and GLUT4 expression in adipocytes (233; 255). One of the best studied inhibitory phosphorylation sites is serine 307/312 (rat/human) of IRS-1. Studies have demonstrated that this residue is phosphorylated by a number of kinases including ERK1/2 (79; 236), JNK (4; 108), IKK β (107), and mammalian target of rapamycin (mTOR) (213).

Transcriptional inhibition by TNF α is accomplished by inhibitory serine phosphorylation of peroxisome proliferator-activated receptor (PPAR) γ by JNK and ERK (2; 184). The induction of insulin resistance and inhibition of PPAR γ by TNF α can be ablated by treatment with thiazolidinediones (TZD) (194; 218; 234). Additionally, TNF α -induced ceramide production is a critical component of insulin resistance (113), leading to inhibition of PKB (269) and decreased CCAAT/enhancer binding protein (C/EBP) α and GLUT4 gene expression (89; 174). Reciprocal positive regulation of both PPAR γ and C/EBP α on each other likely leads to a decrease in mRNA abundance. Because GLUT4 is regulated by both PPAR γ and C/EBP α (127), the decrease in both is likely causal for the decrease of GLUT4 in TNF α induced insulin resistance. Further transcriptional regulation of insulin resistance by TNF α is evidenced by the induction of suppressor of cytokine signaling (SOCS) 3, which blocks IRS-1 tyrosine phosphorylation and thus inhibits insulin-stimulated glucose uptake (78; 274).

Interleukin-6

Like TNF α , IL-6 production is increased in obesity (100; 142; 277) and decreases with weight loss (16). IL-6 signals through the IL-6 receptor which belongs to the class I family of cytokine receptors and utilizes Janus kinases (Jak) as intracellular signaling

pathways leading to activation of signal transducer and activator of transcription (STAT) 1, STAT3, and MAPK (118-119). IL-6 can act both locally and systemically, and approximately 15-35% of circulating IL-6 is adipose tissue derived in humans (195). In human (276), murine (130), and porcine (302) adipocytes IL-6 induces lipolysis. This effect can be inhibited by PPAR γ activation (303). Additionally, IL-6 induces the expression of acute phase proteins (17; 103; 199), a class of proteins secreted predominately by the liver in response to inflammation (i.e TNF α , IL-6).

Literature regarding the ability of IL-6 to induce insulin resistance is much less concise. Similar to TNF α , IL-6 induces the expression of SOCS-3 (85; 152), a key regulatory feature of Jak/STAT cytokine signaling. IL-6 also decreases the expression of the insulin receptor, IRS, GLUT4, PPAR γ , and C/EBP α (152; 232), and these effects are reversed with TZD treatment (152). Unlike TNF α , IL-6 does not induce serine 307 phosphorylation of IRS-1, at least in adipocytes (232). However, evidence suggests that IL-6 signaling can induce inhibitory serine phosphorylation of IRS in other tissues (11; 147; 285).

C-Reactive Protein

CRP is an acute phase protein produced in response to inflammation and tissue damage. CRP is produced predominately by the liver (217), but is also produced by adipocytes (43; 210). Increased expression and circulation of CRP in obesity is likely a response to increased adipose tissue production of IL-6 (308). Increased visceral adipose tissue in obesity likely contributes directly to the increase in circulating concentrations of CRP. Indeed, IL-6 is increased in the portal blood of obese individuals, and is correlated to circulating concentrations of CRP (94). Thus, CRP is highly correlated with obesity, insulin resistance, the metabolic syndrome, and the development of type 2 diabetes (48; 64; 97; 101).

CRP concentrations are negatively correlated with the anti-inflammatory, insulin-sensitizing hormone adiponectin (210) and CRP can induce the expression of MCP-1 (215).

Additionally, mice ubiquitously over-expressing CRP have more adipose tissue macrophages and increased adipose tissue expression of MCP-1 and TNF α (138). These effects were only seen in high fat diet induced obese mice, not lean controls. The increase in adipose tissue macrophages and increased adipose tissue expression of MCP-1 and TNF α in transgenic CRP mice is likely a result of increased activation of macrophages and other immune cells. In addition to inducing MCP-1 expression (215), CRP activation of leukocytes via Fc γ II receptor (CD32) (24) increases CCR2 expression on monocytes (115), the receptor for MCP-1, leading to increased chemotaxis into adipose tissue. Moreover, CRP increases macrophage proliferation (68), by increased macrophage colony-stimulating factor production. Thus, CRP is produced in response to inflammation, but has the potential to further propagate the inflammatory condition.

Monocyte Chemoattractant Protein 1

Recruitment of macrophages into adipose tissue in obesity is largely driven by chemokines, such as MCP-1 (CCL2). In obesity, MCP-1 is elevated (36), both in circulation and in adipose tissue. However, increased adipose tissue expression of MCP-1 is not always reflected in the circulation (61). Increased concentrations of circulating MCP-1 is linked to both insulin resistance (267) and cardiovascular disease (185). Knockout of MCP-1 protects against macrophage infiltration into adipose tissue, insulin resistance, and ectopic lipid deposition (137). Conversely, adipose specific over-expression of MCP-1 increases adipose tissue expression of TNF α and IL-6 (136) and macrophage infiltration (136-137), augments insulin resistance (136-137) and leads to increased ectopic lipid deposition (137). Likewise,

knockout of the MCP-1 receptor, CCR2, in mice results in fewer adipose tissue macrophages and decreased inflammation with reduced insulin resistance and hepatic steatosis (137; 286). The importance of MCP-1 in macrophage infiltration is pivotal to the development of insulin resistance due to the contribution to inflammation by infiltrating macrophages. This is further evidenced by the amelioration of insulin resistance in *db/db* mice expressing a mutant dominant-negative MCP-1 (267). Direct contribution of MCP-1 to insulin resistance is less clear. However, pharmaceutical inhibition of MEK attenuates MCP-1 induced insulin resistance (136), at least in myocytes. Potentially, MCP-1 may contribute directly to insulin resistance in other tissues in addition to increasing macrophage infiltration.

Interleukin-8

Like MCP-1, IL-8 is a chemokine that induces leukocyte extravasation, primarily neutrophils, into tissue during an inflammatory response. IL-8 serum concentrations are increased in obesity and are positively correlated to BMI and negatively associated with insulin sensitivity (144). Additionally, IL-8 is correlated to TNF α concentrations in obese subjects (257). IL-8 can be secreted by a number a cell types, and is secreted by adipocytes (122) after challenge with lipopolysaccharide (LPS) and by adipocytes from insulin resistant subjects (232). The ability of adipocytes to secrete IL-8 suggests that adipose tissue has the potential to recruit neutrophils in obesity due to chronic low-grade inflammation. Indeed, neutrophil infiltration in adipose tissue begins after initial stages of high fat feeding in mice (77), and likely precedes macrophage infiltration. Despite the clear association of IL-8 to obesity, the contribution to insulin resistance remains unclear. Evidence of neutrophil infiltration into adipose tissue following initial stages of high fat feeding disappears after prolonged feeding (77). Additionally, circulating IL-8 concentrations paradoxically

increases after weight loss (38), unlike TNF α and IL-6, and therefore is not associated with improvements in insulin sensitivity following weight loss.

Adiponectin

In addition to inflammatory adipokines, adipose tissue also secretes anti-inflammatory adipokines such as adiponectin (i.e., Acrp30, AdipoQ, apM1). Adiponectin is a 30 kDa protein produced predominantly by adipocytes (180; 239) and circulates in three main forms: trimer, hexamer, and high molecular weight (133). The predominate nuclear receptor for transcriptional control of adiponectin is PPAR γ (126). Additionally, PPAR γ transcriptionally represses the expression of ERp44 (173), which inhibits the secretion of adiponectin from adipocytes (280). Unlike other adipokines, adiponectin is decreased in obesity, insulin resistance, and cardiovascular disease and is negatively correlated with BMI (12; 211). Treatment of human, porcine, and murine macrophages with adiponectin attenuates LPS-induced activation of NF κ B and TNF α production (293; 297; 305). Additionally, adiponectin induces the production of the anti-inflammatory cytokine interleukin-10 (IL-10) in porcine macrophages (293). Similar to macrophages, adiponectin attenuates activation of NF κ B and expression of TNF α and IL-6 in adipocytes (5).

Perturbed glucose homeostasis caused by insulin resistance can be further impaired by decreased concentrations of the anti-diabetic adipokine, adiponectin. Experimental evidence in adiponectin knockout mice suggests that adiponectin is important in protecting against the development of insulin resistance (150; 202). Adiponectin signals through two receptors, AdipoR1 and AdipoR2, which are differentially expressed on multiple cell types (299). Targeted receptor disruption has demonstrated that AdipoR1 and AdipoR2 signaling activates AMP-activated protein kinase (AMPK) and PPAR α (301), respectively. The

insulin-sensitizing effects of adiponectin mediated by activation of AMPK and PPAR α include stimulation of glucose uptake and fatty acid oxidation, and inhibition of lipogenesis and gluconeogenesis (46; 57; 272; 292; 295; 298; 300-301; 306).

Oxidative Stress

Another mechanism of obesity-related insulin resistance includes oxidative stress, which is the generation of reactive oxygen species (ROS) and reactive nitrogen species (RNS). Oxidative stress is closely linked with obesity and type 2 diabetes (13; 19; 26). There are a multitude of ROS and RNS that are generated by NADPH oxidase (20), nitric oxide synthase (214), mitochondrial electron transport (312), and cytochrome P450 (309) in response to stimuli such as TNF α (59), endoplasmic reticulum stress (112), insulin (149), and excess nutrients. Oxidative stress induced with hydrogen peroxide *in vitro* has increased activation of JNK and IKK β , increased serine 307 and 632 phosphorylation of IRS-1, and decreased PI3K activity (25). Additionally, oxidative stress decreases GLUT4 expression in adipocytes (219; 235), thus potentially decreasing insulin stimulated glucose uptake.

Metabolic Endotoxemia

A phenomenon known as metabolic endotoxemia is another contributing factor the development of insulin resistance in obesity. Metabolic endotoxemia is a sub-clinical increase in serum endotoxin, or LPS. LPS initiates a classical immune response via the activation of toll-like receptor 4 (Tlr4), which leads to the production of mediators of inflammation such as TNF α and IL-6. Metabolic endotoxemia is positively correlated with energy intake (8). Consumption of lipids induces postprandial endotoxemia and increases circulating IL-6 (80; 160). Additionally, sub-clinical endotoxemia in humans induces adipose tissue inflammation and transient insulin resistance (190). As such, metabolic

endotoxemia is associated with an increased risk for the development of type 2 diabetes (224).

Obesity and Adipose Tissue Macrophages

Nearly a decade after the link between obesity, inflammation, and insulin resistance was established (123), two laboratories independently discovered that obesity is associated with macrophage accumulation in adipose tissue (287; 296). Both body mass and adipocyte size are positively correlated with the number of macrophages present in adipose tissue (287). In addition to the increase in the number of macrophages, their localization and function change in obese individuals (177). Once within adipose tissue, macrophages recruit more macrophages in a feed forward fashion (206), which increases the inflammatory state and contributes directly to insulin resistance, the metabolic syndrome, and the development of type 2 diabetes.

Human adipose tissue macrophages are characterized via flow cytometry by their expression of CD14, the co-receptor for Tlr4, and CD206, a mannose receptor (310), which distinguishes macrophages from contaminating blood monocytes. Adipose tissue macrophages express high levels of the scavenger receptors CD163 and CD36, do not express the T-cell co-stimulation receptors CD40, CD80, and CD86, and express only low levels of the integrin CD11c (310). One study demonstrated that macrophages expressing CD11c are present in ‘crown-like structures’ surrounding necrotic adipocytes and are involved in lipid uptake (289), consistent with phagocytic activity. Additionally, adipose tissue macrophages produce high concentrations of IL-1 β , IL-6, IL-8, IL-10, TNF α , and MCP-1 (31; 310). Adipose tissue macrophages can also be separated into subpopulations by the expression of CD16, an Fc γ III receptor, with the majority being CD16⁻ (31). However,

the inflammatory gene expression between CD16⁺ and CD16⁻ adipose tissue macrophages differs only by higher expression of MCP-1 and IL-6 in CD16⁻ macrophages (31). The presence of CD16⁻ macrophages in adipose tissue may reflect infiltrating monocytes from the circulation (31), as blood CD16⁻ monocytes express CCR2, the receptor for MCP1 and CD62L (109; 265). However, this has not been definitively proven in regards to adipose tissue macrophages.

Macrophages can be classified into two broad functional subpopulations, which parallels the polarization of naïve CD4⁺ T-lymphocytes toward either a Th1 or Th2 immune response. The first is classically-activated (M1) macrophages. Activation by interferon γ or LPS in a manner consistent with a Th1 immune response characterize the phenotype of M1 macrophages. These cells produce high amounts of TNF α , and express inducible nitric oxide synthase (iNOS). The second is alternatively-activated (M2) macrophages. Activation by IL-4 or IL-13 in a manner consistent with a Th2 immune response characterizes the phenotype of M2 macrophages. These cells produce high amounts of IL-10, and express arginase instead of iNOS. In adipose tissue, resident macrophages display the M2 phenotype. Adipocyte hypertrophy and dysregulation leads to the monocyte diapedesis to adipose tissue. These infiltrated macrophages are of the M1 phenotype. These two subpopulations of macrophages and their functions in adipose tissue are well reviewed by others (50; 62; 198; 243).

From the initial reports in 2003 by Weisberg *et al.* (287) and Xu *et al.* (296), it has been evident macrophages play a pivotal role in the progression of obesity as a state of chronic low-grade inflammation. Much of the research regarding adipose tissue macrophages has been done in mouse models, including macrophage M1 and M2

polarization. While the M1/M2 phenotype is a useful starting point, human macrophage heterogeneity as well as differences between murine and human macrophages emphasizes the need for alternate immunological models to study macrophage function in relation to obesity.

Macrophage infiltration into adipose tissue is largely driven by MCP-1 and its receptor CCR2. CRP is an acute phase protein that interacts with Fc γ II receptor (24) and increases CCR2 expression on monocytes (115), as well as induces the expression of MCP-1 (215). However in mice, CRP is not an acute phase protein and is not highly expressed (264), while it is in humans and swine. The lack of CRP functionality in mice does not abrogate the importance of MCP1 and CCR2 in mice models of obesity, as it is evident that they are intricately involved in macrophage infiltration in mice (136-137; 286), but it clearly demonstrates a difference in the etiology of chronic inflammation in obesity between mice and humans.

Probably most notably is the difference between human and murine macrophages is nitric oxide (NO) production via iNOS. Unlike murine macrophages, human macrophages do not produce NO in response to stimuli (192; 231; 270), which is characteristic of M1 macrophages. Consistent with human macrophages, porcine bone marrow-derived macrophages do not produce NO in response to LPS (139). Additionally, porcine bone marrow-derived macrophages up-regulate genes in response to LPS similar to human macrophages, and that are not up-regulated by murine macrophages. These genes include IDO, STAT4, CCL20 as well others. For further review of similarities and differences between pig and human macrophages, please refer to a review by Fairbairn *et al.* (82).

Recently, phenotypical characterization of Yorkshire swine adipose tissue macrophages was performed by flow cytometry (84). Using the mature macrophage marker

CD203a to identify macrophages, subcutaneous adipose tissue macrophages were predominately CD163⁺, CD16⁺, and CD14⁺ (**Table 1**). It is difficult to make direct comparisons to human adipose tissue macrophages, because the human cells were characterized by being CD14⁺ in conjunction with CD206⁺ (310). Swine adipose tissue macrophages are only 80% CD14⁺ (84). Potentially, the previous studies focusing on human adipose tissue macrophages disregarded populations that are CD14⁻. Although swine adipose tissue macrophages express higher levels of CD16 (**Table 1**) than humans macrophages do (31), they may still be a valuable resource for investigating the function of macrophages in adipose tissue given the greater differences between mice and human macrophages.

Obesity and Fatty Acids

Multiple studies have demonstrated an association between obesity and increased circulating free fatty acids (121; 223; 227). A contributing factor to elevated circulating free fatty acids is increased adipose tissue lipolysis (1; 120; 153; 230), due in part to the decreased anti-lipolytic effect of insulin. The detrimental effects can be evidenced by the induction of insulin resistance in lean human or rats when infused with free fatty acids (164; 166).

The pathological consequences of increased circulating free fatty acids extend to the activation of Tlr4, an event that further propagates chronic inflammation associated with obesity. LPS recognition by Tlr4 is accomplished through the lipid A moiety (225), which contains saturated fatty acids. Early work by Daniel Hwang's laboratory determined that lauric acid induced cyclooxygenase 2 expression in RAW 264.7 macrophages via the activation of Tlr4 and NFκB (163). In adipocytes, treatment with saturated fatty acids induces inflammation including the activation of JNK and NFκB as well as the production of

IL-6 and TNF α (6; 32; 67; 252; 260). Additionally, *in vivo*, the absence or loss-of-function of Tlr4 protects against high saturated fat induced inflammation and insulin resistance (66; 245; 273). There is also evidence that Tlr2 is involved in saturated fatty acid induced inflammation (65; 67; 76)

Beyond Tlr4, saturated fatty acids can induce inflammation and insulin resistance via ceramide production. Lipolysis can induce inflammation via activation of JNK and p38 MAPK pathways. Inhibition of ceramide synthesis blunts this activation (200) in adipocytes. Moreover, inhibition of fatty acid synthase in adipocytes decreases NF κ B activation (6) through the inhibition of ceramide synthesis. Similarly in monocytes, inhibition of ceramide synthesis reduces JNK, ERK, and p38 MAPK activation due to saturated fatty acid treatment (240).

Circulating saturated free fatty acids are positively correlated with circulating IL-6 concentrations (87) and negatively correlated with circulating adiponectin (88). Conversely, n-3 fatty acids are inversely associated with circulating cytokines (90; 134; 193), including CRP, IL-6, and TNF α . The major n-3 fatty acids with biological activity are eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). These fatty acids are readily available from oily marine fish or can be synthesized *in vivo* from α -linolenic acid. However, it is estimated that only 6% of α -linolenic acid may be converted to EPA or DHA *in vivo* (169). The fate of consumed n-3 fatty acids include incorporation into triglycerides and storage in adipose tissue, incorporation into phospholipids to form part of the cellular membrane, circulation as free fatty acids complexed with albumin, or oxidized for energy.

Several studies have demonstrated the immunosuppressive potential of n-3 fatty acids in multiple cell types including T cells, monocytes, macrophages, dendritic cells, and

adipocytes (6; 124; 205; 311; 313). Observational studies demonstrate a beneficial relationship between n-3 fatty acid intake and obesity, but data from cross-sectional studies are limited (39). In rodents, n-3 fatty acids decrease adipose tissue inflammation, macrophage infiltration, and insulin resistance as well as increase circulating adiponectin (91; 125; 135; 271). In swine, EPA and DHA attenuate LPS induced febrile response and circulating TNF α . (104). Molecular mechanisms of inhibition of inflammation by n-3 fatty acids include alterations in lipid rafts (256; 284), altered eicosanoid production (14; 251), inhibition of NF κ B (6; 162; 314), and activation of PPAR γ (15; 207).

Swine Models of Obesity

Swine are increasingly being used as a biomedical research model, in large part due to their similar physiology and anatomy to humans (170). In the context of obesity and metabolic syndrome, swine have been used to study cardiovascular disease (22; 27-29; 33; 58; 75; 106; 110; 154; 188; 196; 201; 203; 216; 229; 261; 294), genetic analysis (83; 95-96; 114; 179; 208; 212; 263; 291), drug testing (42; 226; 304; 315), pancreas function (102; 155; 157), dietary components (34; 221; 228-229; 238), steatohepatitis (21; 165), renal function (47; 168; 171), insulin signaling (161), and exercise training (58; 75).

Commercial swine fed diets high in fat and cholesterol have been used to study diabetes accelerated atherosclerosis (110; 201; 261). These swine developed hypercholesterolemia, but had normal circulating concentrations of glucose and triglycerides. Development of diabetes by treatment with streptozotocin was required to induce hyperglycemia. Additionally, streptozotocin exacerbated the progression of atherosclerosis in this model. Consumption of high fat diets by commercial swine increases abdominal fat, weight gain, and results in hypertension and hypercholesterolemia (106). However, it does

not significantly affect insulin sensitivity or increase circulating concentrations of CRP (106). As mentioned previously, increased circulating free fatty acids and sub-clinical endotoxemia are physiological responses associated with obesity. In commercial swine, experimental endotoxemia and increased circulating concentrations of free fatty acids did not affect glucose homeostasis, but did decrease circulating growth hormone concentrations (40), which may increase adipose tissue mass and decrease lean mass.

Commercial swine have also been used to study the effect of consumption of different fats on body composition and metabolic parameters. Saturated fats (lard), mono-unsaturated fats (olive oil), and sucrose consumption increased whole-body adiposity, but only the mono-unsaturated fat increased visceral adiposity (238). Consumption of either saturated or mono-unsaturated fat, but not sucrose, decreased insulin sensitivity (238). Another study evaluated the effect of the consumption of saturated fats (lard) and polyunsaturated fats (canola oil) on metabolic markers and inflammation. Neither fat source altered circulating IL-6 or CRP concentrations, increased adipose tissue macrophages, or altered insulin sensitivity; although the saturated fat induced hypercholesterolemia (34). Differences in the ability of commercial swine to develop dietary induced changes in insulin sensitivity may be affected by the age that dietary treatment was initiated. Sabin *et al.* (238) began treatment at one week of age, Galili *et al.* (106) began treatment at 3 months of age, and Braucher (34) began treatment at 18 weeks of age. Increased age of initial dietary treatments may reduce the negative effects of dietary fat on insulin sensitivity.

Although commercial swine have been used in obesity-related research, common genetic selection for lean growth limits the potential of this model. Ossabaw swine, however, are a promising model for obesity research, and have already been extensively used

in cardiovascular disease and steatohepatitis research (22; 27-29; 74-75; 154; 165; 203; 216). Ossabaw swine were isolated for ~500 years on Ossabaw Island, just off the coast of Georgia, USA (187). The introduction of new genetics has been limited, but includes one Hampshire boar in the mid 1970s (187). This genetic isolation allowed for the development of a “thrifty genotype” to survive the famine and feast environment. Because of this unique adaptation, Ossabaw swine can store more fat on a kg body weight basis than any other non-domesticated mammal (258).

Ossabaw swine are smaller, slower growing, and have decreased muscle mass compared to domestic swine (41; 60; 182). Increased fat storage and decreased muscle mass in Ossabaw swine may be due to decreased growth hormone secretion in response to insulin (283), but not hyperphagia (281). Ossabaw swine have impaired glucose tolerance (282) and hypertriglyceridemia and hypercholesterolemia (81) as well as a higher capacity for gluconeogenesis, glycolysis, and lipogenesis than lean swine (183).

The aforementioned information regarding Ossabaw swine was collected in the 1970s. Recently, growing concern about the obesity and diabetes epidemic sparked new interest in Ossabaw swine and a breeding colony was established by Michael Sturek (262). Phenotypically, Ossabaw swine display many of the same characteristics today as they did in the 1970’s. Upon development of obesity, they develop hypercholesterolemia (23; 27; 29; 56; 71; 74; 148; 159; 165; 168; 175; 203; 216), hypertriglyceridemia (27; 29; 71; 74; 148; 165; 216), impaired glucose tolerance (74; 159; 203), hyperglycemia (27; 29; 165; 216), and hyperinsulinemia (27; 29; 71; 165). It is important to note that in addition to high fat content, some of the diets contained supra-physiologic amounts of cholesterol (23; 27; 29; 33; 56; 74; 148; 159; 165; 168; 175; 203; 216), which may augment the dyslipidemia in this model.

Only one study showed a decrease in serum adiponectin, marked hyperleptinemia, and systemic inflammation upon prolonged consumption of a high fat diet (165).

Ossabaw swine contain a mutation in the $\gamma 3$ subunit of AMPK (172). This mutation (V224I) is a Val¹⁹⁹→Ile missense mutation that causes low muscle glycogen and increased intramuscular fat (10; 55). This mutation is consistent with the increase in intramuscular lipid seen in obese Ossabaw swine (56). It has been hypothesized by Shen et al. that this mutation may be causal in the development of insulin resistance in Ossabaw swine (244). However, to this date this has not been investigated.

Another common swine model is the Yucatan minipig. Yucatan swine originate from the Yucatan Peninsula of Mexico. When Yucatan swine are fed a high fat high, cholesterol diet, they develop hypercholesterolemia (72; 111; 196; 209; 290), but do not exhibit decreased insulin sensitivity in response to an insulin challenge (209). Yucatan swine do not readily develop hyperglycemia upon consumption of a western diet (30; 72; 196; 209; 241; 290), but do have a decreased glucose infusion rate during the hyperinsulinemic-euglycemic clamp (241). Additionally, Yucatan swine fed a high fat high cholesterol diet have higher liver content of cholesterol, cholesteryl esters, and triglycerides (72), as well as increased visceral adiposity (290). When fed a western diet, Yucatan swine develop hyperleptinemia (241; 290) and have increased adipose tissue expression of leptin and IL-6 (241). Whereas previous studies in Yucatan swine showed only hypercholesterolemia and minimal effect on glucose homeostasis; more recently Yucatan swine have been shown to develop increased circulating concentrations of triglycerides and NEFA, as well as hypercholesterolemia after long-term consumption of a high fat diet (161). In addition, these swine develop hyperglycemia, hyperinsulinemia, and impaired myocardial insulin signaling (161).

Differences between these studies may be attributable to varying genetics between breeding herds.

Göttingen minipigs are another swine model that are used to study obesity and its co-morbidities. Göttingen swine were developed in the 1960s in Göttingen, Germany by cross breeding between Vietnamese swine, Hormel swine, and the German improved Landrace (156). Göttingen swine fed a high fat diet have increased body fat (132; 158; 275), plasma cholesterol (158) and triglycerides (132). Göttingen swine display insulin resistance after 3-5 months of high fat feeding (158; 275), as well as hyperinsulinemia in response to an intravenous glucose tolerance test after as little as 5 weeks of high fat feeding (132). However, impaired fasting glucose was only seen in one study (158). Additionally, there is a sexual dimorphism in the susceptibility of Göttingen swine to negative metabolic effects during diet-induced obesity. Female Göttingen swine become more obese, have a greater degree of insulin resistance, and a more atherogenic plasma profile (i.e. cholesterol, triglycerides) than their male counterparts (52). Higher concentrations of both testosterone and estradiol (52) in male Göttingen swine may reduce their risk for obesity-associated co-morbidities (51).

In addition to Ossabaw, Yucatan, Göttingen, and commercial swine, there are several Chinese swine that have been used in obesity research. These include Guizhou, Ningxiang, Bama, Wuzhishan, and Nongda swine. When fed a high fat, high sucrose diet, Guizhou swine have increased body weight gain and increased fat deposition (294; 304). These swine develop dyslipidemia (294; 304), hyperglycemia (294), and display insulin resistance as evidenced by both an insulin challenge and an oral glucose tolerance test (294; 304). Additionally, ectopic lipid deposition is evident in the liver, pancreas, and muscle after long

term dietary treatment (294; 304). Moreover, Guizhou swine have increased circulating concentrations of TNF α after long-term dietary treatment (294). The genetically obese Chinese Ningxiang swine may have potential as an obesity model. Compared to lean commercial swine Ningxiang have higher circulating concentrations of insulin and cholesterol, as well as greater adipose tissue and less lean mass percentage (117). In a direct comparison of Bama, Wuzhishan, and Nongda swine, only the Bama and Wuzhishan swine developed hyperglycemia in response to a high fat and high sucrose diet (49). Additionally, Bama swine fed a high fat, sucrose, and cholesterol diet developed hyperglycemia, hyperinsulinemia, hypercholesterolemia, and also had decreased circulating concentrations of IGF-1 (167).

While rodent models have been a pillar to obesity research, they do not consistently develop three or more of the risk factors of the metabolic syndrome (253). The ability of Ossabaw and Yucatan swine to develop features of metabolic syndrome has been compared by Neeb *et al.* (203). Yucatan swine only develop hypercholesterolemia, while Ossabaw swine develop obesity, insulin resistance, glucose intolerance, hypertriglyceridemia, hypercholesterolemia, and hypertension (203). Additionally, as reviewed above, Ossabaw swine consistently develop more risk factors for the metabolic syndrome than other swine as breeds. While Ossabaw swine do develop obesity and insulin resistance, adipose tissue inflammation has yet to be evaluated in this breed, or any other breed. Only one study demonstrated increased systemic inflammation in Ossabaw swine (165), but did not evaluate adipose tissue inflammation. Chronic low-grade inflammation is a hallmark of obesity and thus, needs to be evaluated in Ossabaw swine to further the development of the model.

Summary

The prevalence of obesity and its co-morbidities continues to rise in the United States and other developed countries. Excess caloric intake causes adipose tissue expansion through adipocyte hypertrophy and hyperplasia. Adipocyte hypertrophy leads to adipose tissue dysfunction causing increased production of inflammatory adipokines (i.e. IL-6, TNF α , MCP-1) and a decrease in anti-inflammatory adipokines (i.e. adiponectin). These events lead to macrophage infiltration into adipose tissue, propagating inflammation and creating a chronic state of low-grade inflammation that is often associated with obesity. The increased production of inflammatory adipokines by adipocytes and macrophages, as well as oxidative stress induces adipose tissue insulin resistance via the activation of NF κ B and MAPK pathways. Free fatty acids released from adipocytes due to insulin resistance, as well as from the breakdown of necrotic adipocytes by macrophages causes ectopic lipid deposition, particularly in muscle, liver, and pancreas. Ectopic lipid deposition, in conjunction with increased systemic inflammation from adipokines, can lead to insulin resistance in the liver and muscle and decrease pancreatic insulin secretion. These events can lead to the eventual development of type 2 diabetes mellitus.

The susceptibility to obesity is influenced by multiple factors including genotype, environment, and poor diet. Dietary factors, such as fatty acids, have potential implications in obesity. Saturated fatty acids are linked to the inflammatory profile in obese individuals. Saturated fatty acids can activate Tlr4, activating NF κ B and MAPK pathways and thus, increasing inflammation. Additionally, consumption of saturated fatty acids can contribute to metabolic endotoxemia. Conversely, the n-3 fatty acids EPA and DHA are anti-inflammatory in nature and are negatively correlated to the inflammatory profile in obese

individuals. EPA and DHA increase PPAR γ activation and adiponectin production, and decrease NF κ B activation.

Largely because of comparative physiology and anatomy, the use of swine for research involving obesity and its co-morbidities has increased. Ossabaw swine in particular become obese when adequate food is available and develop multiple risk factors of the metabolic syndrome including insulin resistance. However, adipose tissue inflammation has not been investigated in this model. Macrophage infiltration and function in adipose tissue is of particular interest due to immunological discrepancies between mice and human, especially regarding CRP function and NO production by macrophages. Swine are a more relevant immune model for humans, but needs further evaluation. Based upon the literature outlined above, I sought to characterize adipose tissue inflammation in Ossabaw swine in response to saturated and n-3 fatty acids and to determine specific genes involved in adipose tissue expansion by diet-induced obesity to further validate Ossabaw swine as a model for human obesity and obesity-linked inflammation.

References

1. Abdul-Ghani MA, Molina-Carrion M, Jani R, Jenkinson C, Defronzo RA. 2008. Adipocytes in subjects with impaired fasting glucose and impaired glucose tolerance are resistant to the anti-lipolytic effect of insulin. *Acta Diabetol* 45:147-150.
2. Adams M, Reginato MJ, Shao D, Lazar MA, Chatterjee VK. 1997. Transcriptional activation by peroxisome proliferator-activated receptor gamma is inhibited by phosphorylation at a consensus mitogen-activated protein kinase site. *J Biol Chem* 272:5128-5132.
3. Aguilar-Salinas CA, Garcia EG, Robles L, Riano D, Ruiz-Gomez DG, Garcia-Ulloa AC, Melgarejo MA, Zamora M, Guillen-Pineda LE, Mehta R, Canizales-Quinteros S, Tusie Luna MT, Gomez-Perez FJ. 2008. High adiponectin concentrations are associated with the metabolically healthy obese phenotype. *J Clin Endocrinol Metab* 93:4075-4079.

4. Aguirre V, Uchida T, Yenush L, Davis R, White MF. 2000. The c-Jun NH(2)-terminal kinase promotes insulin resistance during association with insulin receptor substrate-1 and phosphorylation of Ser(307). *J Biol Chem* 275:9047-9054.
5. Ajuwon KM, Spurlock ME. 2005. Adiponectin inhibits LPS-induced NF-kappaB activation and IL-6 production and increases PPARgamma2 expression in adipocytes. *Am J Physiol Regul Integr Comp Physiol* 288:R1220-1225.
6. Ajuwon KM, Spurlock ME. 2005. Palmitate activates the NF-kappaB transcription factor and induces IL-6 and TNFalpha expression in 3T3-L1 adipocytes. *J Nutr* 135:1841-1846.
7. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, Fruchart JC, James WP, Loria CM, Smith SC, Jr. 2009. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 120:1640-1645.
8. Amar J, Burcelin R, Ruidavets JB, Cani PD, Fauvel J, Alessi MC, Chamontin B, Ferrieres J. 2008. Energy intake is associated with endotoxemia in apparently healthy men. *Am J Clin Nutr* 87:1219-1223.
9. Andel M, Polak J, Kraml P, Dlouhy P, Stich V. 2009. [Chronic mild inflammation links obesity, metabolic syndrome, atherosclerosis and diabetes]. *Vnitr Lek* 55:659-665.
10. Andersson L. 2003. Identification and characterization of AMPK gamma 3 mutations in the pig. *Biochem Soc Trans* 31:232-235.
11. Andreozzi F, Laratta E, Procopio C, Hribal ML, Sciacqua A, Perticone M, Miele C, Perticone F, Sesti G. 2007. Interleukin-6 impairs the insulin signaling pathway, promoting production of nitric oxide in human umbilical vein endothelial cells. *Mol Cell Biol* 27:2372-2383.
12. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, Hotta K, Shimomura I, Nakamura T, Miyaoka K, Kuriyama H, Nishida M, Yamashita S, Okubo K, Matsubara K, Muraguchi M, Ohmoto Y, Funahashi T, Matsuzawa Y. 1999. Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun* 257:79-83.
13. Atabek ME, Vatansev H, Erkul I. 2004. Oxidative stress in childhood obesity. *J Pediatr Endocrinol Metab* 17:1063-1068.

14. Bagga D, Wang L, Farias-Eisner R, Glaspy JA, Reddy ST. 2003. Differential effects of prostaglandin derived from omega-6 and omega-3 polyunsaturated fatty acids on COX-2 expression and IL-6 secretion. *Proc Natl Acad Sci U S A* 100:1751-1756.
15. Banga A, Unal R, Tripathi P, Pokrovskaya I, Owens RJ, Kern PA, Ranganathan G. 2009. Adiponectin translation is increased by the PPARgamma agonists pioglitazone and omega-3 fatty acids. *Am J Physiol Endocrinol Metab* 296:E480-489.
16. Bastard JP, Jardel C, Bruckert E, Blondy P, Capeau J, Laville M, Vidal H, Hainque B. 2000. Elevated levels of interleukin 6 are reduced in serum and subcutaneous adipose tissue of obese women after weight loss. *J Clin Endocrinol Metab* 85:3338-3342.
17. Bastard JP, Jardel C, Delattre J, Hainque B, Bruckert E, Oberlin F. 1999. Evidence for a link between adipose tissue interleukin-6 content and serum C-reactive protein concentrations in obese subjects. *Circulation* 99:2221-2222.
18. Basu A, Basu R, Shah P, Vella A, Rizza RA, Jensen MD. 2001. Systemic and regional free fatty acid metabolism in type 2 diabetes. *Am J Physiol Endocrinol Metab* 280:E1000-1006.
19. Baynes JW. 1991. Role of oxidative stress in development of complications in diabetes. *Diabetes* 40:405-412.
20. Bedard K, Krause KH. 2007. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* 87:245-313.
21. Bell LN, Lee L, Saxena R, Bemis KG, Wang M, Theodorakis JL, Vuppalanchi R, Alloosh M, Sturek M, Chalasani N. 2010. Serum proteomic analysis of diet-induced steatohepatitis and metabolic syndrome in the Ossabaw miniature swine. *Am J Physiol Gastrointest Liver Physiol* 298:G746-754.
22. Bender SB, Tune JD, Borbouse L, Long X, Sturek M, Laughlin MH. 2009. Altered mechanism of adenosine-induced coronary arteriolar dilation in early-stage metabolic syndrome. *Exp Biol Med (Maywood)* 234:683-692.
23. Berwick ZC, Dick GM, Moberly SP, Kohr MC, Sturek M, Tune JD. 2012. Contribution of voltage-dependent K(+) channels to metabolic control of coronary blood flow. *J Mol Cell Cardiol* 52:912-919.
24. Bharadwaj D, Stein MP, Volzer M, Mold C, Du Clos TW. 1999. The major receptor for C-reactive protein on leukocytes is fcgamma receptor II. *J Exp Med* 190:585-590.

25. Bloch-Damti A, Potashnik R, Gual P, Le Marchand-Brustel Y, Tanti JF, Rudich A, Bashan N. 2006. Differential effects of IRS1 phosphorylated on Ser307 or Ser632 in the induction of insulin resistance by oxidative stress. *Diabetologia* 49:2463-2473.
26. Block G, Dietrich M, Norkus EP, Morrow JD, Hudes M, Caan B, Packer L. 2002. Factors associated with oxidative stress in human populations. *Am J Epidemiol* 156:274-285.
27. Borbouse L, Dick GM, Asano S, Bender SB, Dincer UD, Payne GA, Neeb ZP, Bratz IN, Sturek M, Tune JD. 2009. Impaired function of coronary BK(Ca) channels in metabolic syndrome. *Am J Physiol Heart Circ Physiol* 297:H1629-1637.
28. Borbouse L, Dick GM, Payne GA, Berwick ZC, Neeb ZP, Alloosh M, Bratz IN, Sturek M, Tune JD. 2010. Metabolic syndrome reduces the contribution of K⁺ channels to ischemic coronary vasodilation. *Am J Physiol Heart Circ Physiol* 298:H1182-1189.
29. Borbouse L, Dick GM, Payne GA, Payne BD, Svendsen MC, Neeb ZP, Alloosh M, Bratz IN, Sturek M, Tune JD. 2010. Contribution of BK(Ca) channels to local metabolic coronary vasodilation: Effects of metabolic syndrome. *Am J Physiol Heart Circ Physiol* 298:H966-973.
30. Boullion RD, Mokelke EA, Wamhoff BR, Otis CR, Wenzel J, Dixon JL, Sturek M. 2003. Porcine model of diabetic dyslipidemia: insulin and feed algorithms for mimicking diabetes mellitus in humans. *Comp Med* 53:42-52.
31. Bourlier V, Zakaroff-Girard A, Miranville A, De Barros S, Maumus M, Sengenès C, Galitzky J, Lafontan M, Karpe F, Frayn KN, Bouloumie A. 2008. Remodeling phenotype of human subcutaneous adipose tissue macrophages. *Circulation* 117:806-815.
32. Bradley RL, Fisher FF, Maratos-Flier E. 2008. Dietary fatty acids differentially regulate production of TNF- α and IL-10 by murine 3T3-L1 adipocytes. *Obesity (Silver Spring)* 16:938-944.
33. Bratz IN, Dick GM, Tune JD, Edwards JM, Neeb ZP, Dincer UD, Sturek M. 2008. Impaired capsaicin-induced relaxation of coronary arteries in a porcine model of the metabolic syndrome. *Am J Physiol Heart Circ Physiol* 294:H2489-2496.
34. Braucher DR. The role of adiponectin on body composition, adipocyte differentiation, preadipocyte proliferation and its modulation of inflammation with high fat diets. *Food Science and Human Nutrition*. Ames: Iowa State University.
35. Brochu M, Tchernof A, Dionne IJ, Sites CK, Eltabbakh GH, Sims EA, Poehlman ET. 2001. What are the physical characteristics associated with a normal metabolic profile

- despite a high level of obesity in postmenopausal women? *J Clin Endocrinol Metab* 86:1020-1025.
36. Bruun JM, Lihn AS, Pedersen SB, Richelsen B. 2005. Monocyte chemoattractant protein-1 release is higher in visceral than subcutaneous human adipose tissue (AT): implication of macrophages resident in the AT. *J Clin Endocrinol Metab* 90:2282-2289.
 37. Bruun JM, Lihn AS, Verdich C, Pedersen SB, Toubro S, Astrup A, Richelsen B. 2003. Regulation of adiponectin by adipose tissue-derived cytokines: in vivo and in vitro investigations in humans. *Am J Physiol Endocrinol Metab* 285:E527-533.
 38. Bruun JM, Verdich C, Toubro S, Astrup A, Richelsen B. 2003. Association between measures of insulin sensitivity and circulating levels of interleukin-8, interleukin-6 and tumor necrosis factor-alpha. Effect of weight loss in obese men. *Eur J Endocrinol* 148:535-542.
 39. Buckley JD, Howe PR. 2010. Long-chain omega-3 polyunsaturated fatty acids may be beneficial for reducing obesity-a review. *Nutrients* 2:1212-1230.
 40. Buhl M, Gjedsted J, Granfeldt A, Larsen PO, Schmitz O, Tonnesen E, Moller N. 2010. Metabolic effects of free fatty acids during endotoxaemia in a porcine model--free fatty acid inhibition of growth hormone secretion as a potential catabolic feedback mechanism. *Horm Metab Res* 42:348-352.
 41. Buhlinger CA, Wangsness PJ, Martin RJ, Ziegler JH. 1978. Body composition, in vitro lipid metabolism and skeletal muscle characteristics in fast-growing, lean and in slow-growing, obese pigs at equal age and weight. *Growth* 42:225-236.
 42. Cai M, Yin W, Li Q, Liao D, Tsutsumi K, Hou H, Liu Y, Zhang C, Li J, Wang Z, Xiao J. 2006. Effects of NO-1886 on inflammation-associated cytokines in high-fat/high-sucrose/high-cholesterol diet-fed miniature pigs. *Eur J Pharmacol* 540:139-146.
 43. Calabro P, Chang DW, Willerson JT, Yeh ET. 2005. Release of C-reactive protein in response to inflammatory cytokines by human adipocytes: linking obesity to vascular inflammation. *J Am Coll Cardiol* 46:1112-1113.
 44. Cauchi S, Stutzmann F, Cavalcanti-Proenca C, Durand E, Pouta A, Hartikainen AL, Marre M, Vol S, Tammelin T, Laitinen J, Gonzalez-Izquierdo A, Blakemore AI, Elliott P, Meyre D, Balkau B, Jarvelin MR, Froguel P. 2009. Combined effects of MC4R and FTO common genetic variants on obesity in European general populations. *J Mol Med (Berl)* 87:537-546.

45. Cawthorn WP, Sethi JK. 2008. TNF-alpha and adipocyte biology. *FEBS Lett* 582:117-131.
46. Ceddia RB, Somwar R, Maida A, Fang X, Bikopoulos G, Sweeney G. 2005. Globular adiponectin increases GLUT4 translocation and glucose uptake but reduces glycogen synthesis in rat skeletal muscle cells. *Diabetologia* 48:132-139.
47. Chade AR, Bentley MD, Zhu X, Rodriguez-Porcel M, Niemeyer S, Amores-Arriaga B, Napoli C, Ritman EL, Lerman A, Lerman LO. 2004. Antioxidant intervention prevents renal neovascularization in hypercholesterolemic pigs. *J Am Soc Nephrol* 15:1816-1825.
48. Chambers JC, Eda S, Bassett P, Karim Y, Thompson SG, Gallimore JR, Pepys MB, Kooner JS. 2001. C-reactive protein, insulin resistance, central obesity, and coronary heart disease risk in Indian Asians from the United Kingdom compared with European whites. *Circulation* 104:145-150.
49. Chen H, Liu YQ, Li CH, Guo XM, Huang LJ. 2009. The susceptibility of three strains of Chinese minipigs to diet-induced type 2 diabetes mellitus. *Lab Anim (NY)* 38:355-363.
50. Chinetti-Gbaguidi G, Staels B. 2011. Macrophage polarization in metabolic disorders: functions and regulation. *Curr Opin Lipidol* 22:365-372.
51. Christoffersen BO, Gade LP, Golozoubova V, Svendsen O, Raun K. 2010. Influence of castration-induced testosterone and estradiol deficiency on obesity and glucose metabolism in male Gottingen minipigs. *Steroids* 75:676-684.
52. Christoffersen BO, Grand N, Golozoubova V, Svendsen O, Raun K. 2007. Gender-associated differences in metabolic syndrome-related parameters in Gottingen minipigs. *Comp Med* 57:493-504.
53. Cinti S. 2005. The adipose organ. *Prostaglandins Leukot Essent Fatty Acids* 73:9-15.
54. Cinti S, Mitchell G, Barbatelli G, Murano I, Ceresi E, Faloia E, Wang S, Fortier M, Greenberg AS, Obin MS. 2005. Adipocyte death defines macrophage localization and function in adipose tissue of obese mice and humans. *J Lipid Res* 46:2347-2355.
55. Ciobanu D, Bastiaansen J, Malek M, Helm J, Woollard J, Plastow G, Rothschild M. 2001. Evidence for new alleles in the protein kinase adenosine monophosphate-activated gamma(3)-subunit gene associated with low glycogen content in pig skeletal muscle and improved meat quality. *Genetics* 159:1151-1162.

56. Clark BA, Alloosh M, Wenzel JW, Sturek M, Kostrominova TY. 2011. Effect of diet-induced obesity and metabolic syndrome on skeletal muscles of Ossabaw miniature swine. *Am J Physiol Endocrinol Metab* 300:E848-857.
57. Combs TP, Berg AH, Obici S, Scherer PE, Rossetti L. 2001. Endogenous glucose production is inhibited by the adipose-derived protein Acrp30. *J Clin Invest* 108:1875-1881.
58. Company JM, Booth FW, Laughlin MH, Arce-Esquivel AA, Sacks HS, Bahouth SW, Fain JN. 2010. Epicardial fat gene expression after aerobic exercise training in pigs with coronary atherosclerosis: relationship to visceral and subcutaneous fat. *J Appl Physiol* 109:1904-1912.
59. Corda S, Laplace C, Vicaute E, Duranteau J. 2001. Rapid reactive oxygen species production by mitochondria in endothelial cells exposed to tumor necrosis factor- α is mediated by ceramide. *Am J Respir Cell Mol Biol* 24:762-768.
60. Cote PJ, Wangsness PJ. 1978. Rate, Composition and Efficiency of Growth in Lean and Obese Pigs. *J Anim Sci* 47:441-447.
61. Dahlman I, Kaaman M, Olsson T, Tan GD, Bickerton AS, Wahlen K, Andersson J, Nordstrom EA, Blomqvist L, Sjogren A, Forsgren M, Attersand A, Arner P. 2005. A unique role of monocyte chemoattractant protein 1 among chemokines in adipose tissue of obese subjects. *J Clin Endocrinol Metab* 90:5834-5840.
62. Dalmas E, Clement K, Guerre-Millo M. 2011. Defining macrophage phenotype and function in adipose tissue. *Trends Immunol* 32:307-314.
63. Dandona P, Weinstock R, Thusu K, Abdel-Rahman E, Aljada A, Wadden T. 1998. Tumor necrosis factor- α in sera of obese patients: fall with weight loss. *J Clin Endocrinol Metab* 83:2907-2910.
64. Danesh J, Muir J, Wong YK, Ward M, Gallimore JR, Pepys MB. 1999. Risk factors for coronary heart disease and acute-phase proteins. A population-based study. *Eur Heart J* 20:954-959.
65. Davis JE, Braucher DR, Walker-Daniels J, Spurlock ME. 2011. Absence of Tlr2 protects against high-fat diet-induced inflammation and results in greater insulin-stimulated glucose transport in cultured adipocytes. *J Nutr Biochem* 22:136-141.
66. Davis JE, Gabler NK, Walker-Daniels J, Spurlock ME. 2008. Tlr-4 deficiency selectively protects against obesity induced by diets high in saturated fat. *Obesity (Silver Spring)* 16:1248-1255.

67. Davis JE, Gabler NK, Walker-Daniels J, Spurlock ME. 2009. The c-Jun N-terminal kinase mediates the induction of oxidative stress and insulin resistance by palmitate and toll-like receptor 2 and 4 ligands in 3T3-L1 adipocytes. *Horm Metab Res* 41:523-530.
68. Devaraj S, Yun JM, Duncan-Staley C, Jialal I. 2009. C-reactive protein induces M-CSF release and macrophage proliferation. *J Leukoc Biol* 85:262-267.
69. DiGirolamo M, Fine JB, Tagra K, Rossmanith R. 1998. Qualitative regional differences in adipose tissue growth and cellularity in male Wistar rats fed ad libitum. *Am J Physiol* 274:R1460-1467.
70. Dina C, Meyre D, Gallina S, Durand E, Korner A, Jacobson P, Carlsson LM, Kiess W, Vatin V, Lecoecur C, Delplanque J, Vaillant E, Pattou F, Ruiz J, Weill J, Levy-Marchal C, Horber F, Potoczna N, Hercberg S, Le Stunff C, Bougneres P, Kovacs P, Marre M, Balkau B, Cauchi S, Chevre JC, Froguel P. 2007. Variation in FTO contributes to childhood obesity and severe adult obesity. *Nat Genet* 39:724-726.
71. Dincer UD. 2011. Cardiac beta-adrenoceptor expression is markedly depressed in Ossabaw swine model of cardiometabolic risk. *Int J Gen Med* 4:493-499.
72. Dixon JL, Shen S, Vuchetich JP, Wysocka E, Sun GY, Sturek M. 2002. Increased atherosclerosis in diabetic dyslipidemic swine: protection by atorvastatin involves decreased VLDL triglycerides but minimal effects on the lipoprotein profile. *J Lipid Res* 43:1618-1629.
73. Donnelly KL, Smith CI, Schwarzenberg SJ, Jessurun J, Boldt MD, Parks EJ. 2005. Sources of fatty acids stored in liver and secreted via lipoproteins in patients with nonalcoholic fatty liver disease. *J Clin Invest* 115:1343-1351.
74. Dyson MC, Alloosh M, Vuchetich JP, Mokelke EA, Sturek M. 2006. Components of metabolic syndrome and coronary artery disease in female Ossabaw swine fed excess atherogenic diet. *Comp Med* 56:35-45.
75. Edwards JM, Neeb ZP, Alloosh MA, Long X, Bratz IN, Peller CR, Byrd JP, Kumar S, Obukhov AG, Sturek M. 2010. Exercise training decreases store-operated Ca²⁺ entry associated with metabolic syndrome and coronary atherosclerosis. *Cardiovasc Res* 85:631-640.
76. Ehses JA, Meier DT, Wueest S, Rytka J, Boller S, Wielinga PY, Schraenen A, Lemaire K, Debray S, Van Lommel L, Pospisilik JA, Tschopp O, Schultze SM, Malipiero U, Esterbauer H, Ellingsgaard H, Rutti S, Schuit FC, Lutz TA, Boni-Schnetzler M, Konrad D, Donath MY. 2010. Toll-like receptor 2-deficient mice are protected from insulin resistance and beta cell dysfunction induced by a high-fat diet. *Diabetologia* 53:1795-1806.

77. Elgazar-Carmon V, Rudich A, Hadad N, Levy R. 2008. Neutrophils transiently infiltrate intra-abdominal fat early in the course of high-fat feeding. *J Lipid Res* 49:1894-1903.
78. Emanuelli B, Peraldi P, Filloux C, Chavey C, Freidinger K, Hilton DJ, Hotamisligil GS, Van Obberghen E. 2001. SOCS-3 inhibits insulin signaling and is up-regulated in response to tumor necrosis factor-alpha in the adipose tissue of obese mice. *J Biol Chem* 276:47944-47949.
79. Engelman JA, Berg AH, Lewis RY, Lisanti MP, Scherer PE. 2000. Tumor necrosis factor alpha-mediated insulin resistance, but not dedifferentiation, is abrogated by MEK1/2 inhibitors in 3T3-L1 adipocytes. *Mol Endocrinol* 14:1557-1569.
80. Erridge C, Attina T, Spickett CM, Webb DJ. 2007. A high-fat meal induces low-grade endotoxemia: evidence of a novel mechanism of postprandial inflammation. *Am J Clin Nutr* 86:1286-1292.
81. Etherton TD, Kris-Etherton PM. 1980. Characterization of plasma lipoproteins in swine with different propensities for obesity. *Lipids* 15:823-829.
82. Fairbairn L, Kapetanovic R, Sester DP, Hume DA. 2011. The mononuclear phagocyte system of the pig as a model for understanding human innate immunity and disease. *J Leukoc Biol* 89:855-871.
83. Fan B, Du ZQ, Rothschild MF. 2009. The fat mass and obesity-associated (FTO) gene is associated with intramuscular fat content and growth rate in the pig. *Anim Biotechnol* 20:58-70.
84. Faris RJ, Walker-Daniels J, Jones DE, Spurlock ME. 2012. Characterization of swine adipose tissue monocytic lineage cells. *The FASEB Journal* 26:819.842.
85. Fasshauer M, Kralisch S, Klier M, Lossner U, Bluher M, Klein J, Paschke R. 2004. Insulin resistance-inducing cytokines differentially regulate SOCS mRNA expression via growth factor- and Jak/Stat-signaling pathways in 3T3-L1 adipocytes. *J Endocrinol* 181:129-138.
86. Feinstein R, Kanety H, Papa MZ, Lunenfeld B, Karasik A. 1993. Tumor necrosis factor-alpha suppresses insulin-induced tyrosine phosphorylation of insulin receptor and its substrates. *J Biol Chem* 268:26055-26058.
87. Fernandez-Real JM, Broch M, Vendrell J, Ricart W. 2003. Insulin resistance, inflammation, and serum fatty acid composition. *Diabetes Care* 26:1362-1368.

88. Fernandez-Real JM, Vendrell J, Ricart W. 2005. Circulating adiponectin and plasma fatty acid profile. *Clin Chem* 51:603-609.
89. Fernandez-Veledo S, Hernandez R, Teruel T, Mas JA, Ros M, Lorenzo M. 2006. Ceramide mediates TNF-alpha-induced insulin resistance on GLUT4 gene expression in brown adipocytes. *Arch Physiol Biochem* 112:13-22.
90. Ferrucci L, Cherubini A, Bandinelli S, Bartali B, Corsi A, Lauretani F, Martin A, Andres-Lacueva C, Senin U, Guralnik JM. 2006. Relationship of plasma polyunsaturated fatty acids to circulating inflammatory markers. *J Clin Endocrinol Metab* 91:439-446.
91. Flachs P, Mohamed-Ali V, Horakova O, Rossmeisl M, Hosseinzadeh-Attar MJ, Hensler M, Ruzickova J, Kopecky J. 2006. Polyunsaturated fatty acids of marine origin induce adiponectin in mice fed a high-fat diet. *Diabetologia* 49:394-397.
92. Flegal KM, Carroll MD, Ogden CL, Curtin LR. 2010. Prevalence and trends in obesity among US adults, 1999-2008. *JAMA* 303:235-241.
93. Flegal KM, Carroll MD, Ogden CL, Johnson CL. 2002. Prevalence and trends in obesity among US adults, 1999-2000. *JAMA* 288:1723-1727.
94. Fontana L, Eagon JC, Trujillo ME, Scherer PE, Klein S. 2007. Visceral fat adipokine secretion is associated with systemic inflammation in obese humans. *Diabetes* 56:1010-1013.
95. Fontanesi L, Scotti E, Buttazzoni L, Dall'Olio S, Bagnato A, Lo Fiego DP, Davoli R, Russo V. 2010. Confirmed association between a single nucleotide polymorphism in the FTO gene and obesity-related traits in heavy pigs. *Mol Biol Rep* 37:461-466.
96. Fontanesi L, Scotti E, Buttazzoni L, Davoli R, Russo V. 2009. The porcine fat mass and obesity associated (FTO) gene is associated with fat deposition in Italian Duroc pigs. *Anim Genet* 40:90-93.
97. Ford ES. 1999. Body mass index, diabetes, and C-reactive protein among U.S. adults. *Diabetes Care* 22:1971-1977.
98. Ford ES, Schulze MB, Pischon T, Bergmann MM, Joost HG, Boeing H. 2008. Metabolic syndrome and risk of incident diabetes: findings from the European Prospective Investigation into Cancer and Nutrition-Potsdam Study. *Cardiovasc Diabetol* 7:35.
99. Frayling TM, Timpson NJ, Weedon MN, Zeggini E, Freathy RM, Lindgren CM, Perry JR, Elliott KS, Lango H, Rayner NW, Shields B, Harries LW, Barrett JC, Ellard S, Groves CJ, Knight B, Patch AM, Ness AR, Ebrahim S, Lawlor DA, Ring SM, Ben-Shlomo Y, Jarvelin MR, Sovio U, Bennett AJ, Melzer D, Ferrucci L, Loos RJ,

- Barroso I, Wareham NJ, Karpe F, Owen KR, Cardon LR, Walker M, Hitman GA, Palmer CN, Doney AS, Morris AD, Smith GD, Hattersley AT, McCarthy MI. 2007. A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* 316:889-894.
100. Fried SK, Bunkin DA, Greenberg AS. 1998. Omental and subcutaneous adipose tissues of obese subjects release interleukin-6: depot difference and regulation by glucocorticoid. *J Clin Endocrinol Metab* 83:847-850.
 101. Frohlich M, Imhof A, Berg G, Hutchinson WL, Pepys MB, Boeing H, Muche R, Brenner H, Koenig W. 2000. Association between C-reactive protein and features of the metabolic syndrome: a population-based study. *Diabetes Care* 23:1835-1839.
 102. Fullenkamp AM, Bell LN, Robbins RD, Lee L, Saxena R, Alloosh M, Klaunig JE, Mirmira RG, Sturek M, Chalasan N. 2011. Effect of different obesogenic diets on pancreatic histology in Ossabaw miniature swine. *Pancreas* 40:438-443.
 103. Gabay C, Kushner I. 1999. Acute-phase proteins and other systemic responses to inflammation. *N Engl J Med* 340:448-454.
 104. Gabler NK, Spencer JD, Webel DM, Spurlock ME. 2008. n-3 PUFA attenuate lipopolysaccharide-induced down-regulation of toll-like receptor 4 expression in porcine adipose tissue but does not alter the expression of other immune modulators. *J Nutr Biochem* 19:8-15.
 105. Galassi A, Reynolds K, He J. 2006. Metabolic syndrome and risk of cardiovascular disease: a meta-analysis. *Am J Med* 119:812-819.
 106. Galili O, Versari D, Sattler KJ, Olson ML, Mannheim D, McConnell JP, Chade AR, Lerman LO, Lerman A. 2007. Early experimental obesity is associated with coronary endothelial dysfunction and oxidative stress. *Am J Physiol Heart Circ Physiol* 292:H904-911.
 107. Gao Z, Hwang D, Bataille F, Lefevre M, York D, Quon MJ, Ye J. 2002. Serine phosphorylation of insulin receptor substrate 1 by inhibitor kappa B kinase complex. *J Biol Chem* 277:48115-48121.
 108. Gao Z, Zuberi A, Quon MJ, Dong Z, Ye J. 2003. Aspirin inhibits serine phosphorylation of insulin receptor substrate 1 in tumor necrosis factor-treated cells through targeting multiple serine kinases. *J Biol Chem* 278:24944-24950.
 109. Geissmann F, Jung S, Littman DR. 2003. Blood monocytes consist of two principal subsets with distinct migratory properties. *Immunity* 19:71-82.

110. Gerrity RG, Natarajan R, Nadler JL, Kimsey T. 2001. Diabetes-induced accelerated atherosclerosis in swine. *Diabetes* 50:1654-1665.
111. Glazer HP, Osipov RM, Clements RT, Sellke FW, Bianchi C. 2009. Hypercholesterolemia is associated with hyperactive cardiac mTORC1 and mTORC2 signaling. *Cell Cycle* 8:1738-1746.
112. Gregor MF, Hotamisligil GS. 2007. Thematic review series: Adipocyte Biology. Adipocyte stress: the endoplasmic reticulum and metabolic disease. *J Lipid Res* 48:1905-1914.
113. Grigsby RJ, Dobrowsky RT. 2001. Inhibition of ceramide production reverses TNF-induced insulin resistance. *Biochem Biophys Res Commun* 287:1121-1124.
114. Guyonnet-Duperat V, Geverink N, Plastow GS, Evans G, Ousova O, Croisietiere C, Foury A, Richard E, Mormede P, Moisan MP. 2006. Functional implication of an Arg307Gly substitution in corticosteroid-binding globulin, a candidate gene for a quantitative trait locus associated with cortisol variability and obesity in pig. *Genetics* 173:2143-2149.
115. Han KH, Hong KH, Park JH, Ko J, Kang DH, Choi KJ, Hong MK, Park SW, Park SJ. 2004. C-reactive protein promotes monocyte chemoattractant protein-1--mediated chemotaxis through upregulating CC chemokine receptor 2 expression in human monocytes. *Circulation* 109:2566-2571.
116. Hausman DB, DiGirolamo M, Bartness TJ, Hausman GJ, Martin RJ. 2001. The biology of white adipocyte proliferation. *Obes Rev* 2:239-254.
117. He Q, Ren P, Kong X, Wu Y, Wu G, Li P, Hao F, Tang H, Blachier F, Yin Y. 2012. Comparison of serum metabolite compositions between obese and lean growing pigs using an NMR-based metabonomic approach. *J Nutr Biochem* 23:133-139.
118. Heinrich PC, Behrmann I, Haan S, Hermanns HM, Muller-Newen G, Schaper F. 2003. Principles of interleukin (IL)-6-type cytokine signalling and its regulation. *Biochem J* 374:1-20.
119. Heinrich PC, Behrmann I, Muller-Newen G, Schaper F, Graeve L. 1998. Interleukin-6-type cytokine signalling through the gp130/Jak/STAT pathway. *Biochem J* 334 (Pt 2):297-314.
120. Hellstrom L, Reynisdottir S, Langin D, Rossner S, Arner P. 1996. Regulation of lipolysis in fat cells of obese women during long-term hypocaloric diet. *Int J Obes Relat Metab Disord* 20:745-752.

121. Heptulla R, Smitten A, Teague B, Tamborlane WV, Ma YZ, Caprio S. 2001. Temporal patterns of circulating leptin levels in lean and obese adolescents: relationships to insulin, growth hormone, and free fatty acids rhythmicity. *J Clin Endocrinol Metab* 86:90-96.
122. Hoch M, Eberle AN, Peterli R, Peters T, Seboek D, Keller U, Muller B, Linscheid P. 2008. LPS induces interleukin-6 and interleukin-8 but not tumor necrosis factor-alpha in human adipocytes. *Cytokine* 41:29-37.
123. Hotamisligil GS, Shargill NS, Spiegelman BM. 1993. Adipose expression of tumor necrosis factor-alpha: direct role in obesity-linked insulin resistance. *Science* 259:87-91.
124. Hughes DA, Pinder AC. 2000. n-3 polyunsaturated fatty acids inhibit the antigen-presenting function of human monocytes. *Am J Clin Nutr* 71:357S-360S.
125. Itoh M, Suganami T, Satoh N, Tanimoto-Koyama K, Yuan X, Tanaka M, Kawano H, Yano T, Aoe S, Takeya M, Shimatsu A, Kuzuya H, Kamei Y, Ogawa Y. 2007. Increased adiponectin secretion by highly purified eicosapentaenoic acid in rodent models of obesity and human obese subjects. *Arterioscler Thromb Vasc Biol* 27:1918-1925.
126. Iwaki M, Matsuda M, Maeda N, Funahashi T, Matsuzawa Y, Makishima M, Shimomura I. 2003. Induction of adiponectin, a fat-derived antidiabetic and antiatherogenic factor, by nuclear receptors. *Diabetes* 52:1655-1663.
127. Jain RG, Phelps KD, Pekala PH. 1999. Tumor necrosis factor-alpha initiated signal transduction in 3T3-L1 adipocytes. *J Cell Physiol* 179:58-66.
128. Jennings CL, Lambert EV, Collins M, Joffe Y, Levitt NS, Goedecke JH. 2008. Determinants of insulin-resistant phenotypes in normal-weight and obese Black African women. *Obesity (Silver Spring)* 16:1602-1609.
129. Jernas M, Palming J, Sjöholm K, Jennische E, Svensson PA, Gabrielsson BG, Levin M, Sjögren A, Rudemo M, Lystig TC, Carlsson B, Carlsson LM, Lonn M. 2006. Separation of human adipocytes by size: hypertrophic fat cells display distinct gene expression. *Faseb J* 20:1540-1542.
130. Ji C, Chen X, Gao C, Jiao L, Wang J, Xu G, Fu H, Guo X, Zhao Y. 2011. IL-6 induces lipolysis and mitochondrial dysfunction, but does not affect insulin-mediated glucose transport in 3T3-L1 adipocytes. *J Bioenerg Biomembr* 43:367-375.
131. Jiao P, Chen Q, Shah S, Du J, Tao B, Tzamelis I, Yan W, Xu H. 2009. Obesity-related upregulation of monocyte chemotactic factors in adipocytes: involvement of nuclear factor-kappaB and c-Jun NH2-terminal kinase pathways. *Diabetes* 58:104-115.

132. Johansen T, Hansen HS, Richelsen B, Malmlof R. 2001. The obese Gottingen minipig as a model of the metabolic syndrome: dietary effects on obesity, insulin sensitivity, and growth hormone profile. *Comp Med* 51:150-155.
133. Kadowaki T, Yamauchi T, Kubota N, Hara K, Ueki K, Tobe K. 2006. Adiponectin and adiponectin receptors in insulin resistance, diabetes, and the metabolic syndrome. *J Clin Invest* 116:1784-1792.
134. Kalogeropoulos N, Panagiotakos DB, Pitsavos C, Chrysoshoou C, Rousinou G, Toutouza M, Stefanadis C. 2010. Unsaturated fatty acids are inversely associated and n-6/n-3 ratios are positively related to inflammation and coagulation markers in plasma of apparently healthy adults. *Clin Chim Acta* 411:584-591.
135. Kalupahana NS, Claycombe K, Newman SJ, Stewart T, Siriwardhana N, Matthan N, Lichtenstein AH, Moustaid-Moussa N. 2010. Eicosapentaenoic acid prevents and reverses insulin resistance in high-fat diet-induced obese mice via modulation of adipose tissue inflammation. *J Nutr* 140:1915-1922.
136. Kamei N, Tobe K, Suzuki R, Ohsugi M, Watanabe T, Kubota N, Ohtsuka-Kowatari N, Kumagai K, Sakamoto K, Kobayashi M, Yamauchi T, Ueki K, Oishi Y, Nishimura S, Manabe I, Hashimoto H, Ohnishi Y, Ogata H, Tokuyama K, Tsunoda M, Ide T, Murakami K, Nagai R, Kadowaki T. 2006. Overexpression of monocyte chemoattractant protein-1 in adipose tissues causes macrophage recruitment and insulin resistance. *J Biol Chem* 281:26602-26614.
137. Kanda H, Tateya S, Tamori Y, Kotani K, Hiasa K, Kitazawa R, Kitazawa S, Miyachi H, Maeda S, Egashira K, Kasuga M. 2006. MCP-1 contributes to macrophage infiltration into adipose tissue, insulin resistance, and hepatic steatosis in obesity. *J Clin Invest* 116:1494-1505.
138. Kaneko H, Anzai T, Nagai T, Anzai A, Takahashi T, Mano Y, Morimoto K, Maekawa Y, Itoh H, Yoshikawa T, Ogawa S, Fukuda K. 2011. Human C-reactive protein exacerbates metabolic disorders in association with adipose tissue remodelling. *Cardiovasc Res* 91:546-555.
139. Kapetanovic R, Fairbairn L, Beraldi D, Sester DP, Archibald AL, Tuggle CK, Hume DA. 2012. Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. *J Immunol* 188:3382-3394.
140. Karelis AD, Faraj M, Bastard JP, St-Pierre DH, Brochu M, Prud'homme D, Rabasa-Lhoret R. 2005. The metabolically healthy but obese individual presents a favorable inflammation profile. *J Clin Endocrinol Metab* 90:4145-4150.

141. Karelis AD, St-Pierre DH, Conus F, Rabasa-Lhoret R, Poehlman ET. 2004. Metabolic and body composition factors in subgroups of obesity: what do we know? *J Clin Endocrinol Metab* 89:2569-2575.
142. Kern PA, Ranganathan S, Li C, Wood L, Ranganathan G. 2001. Adipose tissue tumor necrosis factor and interleukin-6 expression in human obesity and insulin resistance. *Am J Physiol Endocrinol Metab* 280:E745-751.
143. Kern PA, Saghizadeh M, Ong JM, Bosch RJ, Deem R, Simsolo RB. 1995. The expression of tumor necrosis factor in human adipose tissue. Regulation by obesity, weight loss, and relationship to lipoprotein lipase. *J Clin Invest* 95:2111-2119.
144. Kim CS, Park HS, Kawada T, Kim JH, Lim D, Hubbard NE, Kwon BS, Erickson KL, Yu R. 2006. Circulating levels of MCP-1 and IL-8 are elevated in human obese subjects and associated with obesity-related parameters. *Int J Obes (Lond)* 30:1347-1355.
145. Kim JY, van de Wall E, Laplante M, Azzara A, Trujillo ME, Hofmann SM, Schraw T, Durand JL, Li H, Li G, Jelicks LA, Mehler MF, Hui DY, Deshaies Y, Shulman GI, Schwartz GJ, Scherer PE. 2007. Obesity-associated improvements in metabolic profile through expansion of adipose tissue. *J Clin Invest* 117:2621-2637.
146. Kim LJ, Nalls MA, Eiriksdottir G, Sigurdsson S, Launer LJ, Koster A, Chaves PH, Jonsdottir B, Garcia M, Gudnason V, Harris TB. 2011. Associations of visceral and liver fat with the metabolic syndrome across the spectrum of obesity: the AGES-Reykjavik study. *Obesity (Silver Spring)* 19:1265-1271.
147. Klover PJ, Zimmers TA, Koniaris LG, Mooney RA. 2003. Chronic exposure to interleukin-6 causes hepatic insulin resistance in mice. *Diabetes* 52:2784-2789.
148. Kreutz RP, Alloosh M, Mansour K, Neeb Z, Kreutz Y, Flockhart DA, Sturek M. 2011. Morbid obesity and metabolic syndrome in Ossabaw miniature swine are associated with increased platelet reactivity. *Diabetes Metab Syndr Obes* 4:99-105.
149. Krieger-Brauer HI, Kather H. 1992. Human fat cells possess a plasma membrane-bound H₂O₂-generating system that is activated by insulin via a mechanism bypassing the receptor kinase. *J Clin Invest* 89:1006-1013.
150. Kubota N, Terauchi Y, Yamauchi T, Kubota T, Moroi M, Matsui J, Eto K, Yamashita T, Kamon J, Satoh H, Yano W, Froguel P, Nagai R, Kimura S, Kadowaki T, Noda T. 2002. Disruption of adiponectin causes insulin resistance and neointimal formation. *J Biol Chem* 277:25863-25866.
151. Kuhlmann J, Neumann-Haefelin C, Belz U, Kalisch J, Juretschke HP, Stein M, Kleinschmidt E, Kramer W, Herling AW. 2003. Intramyocellular lipid and insulin

- resistance: a longitudinal in vivo ¹H-spectroscopic study in Zucker diabetic fatty rats. *Diabetes* 52:138-144.
152. Lagathu C, Bastard JP, Auclair M, Maachi M, Capeau J, Caron M. 2003. Chronic interleukin-6 (IL-6) treatment increased IL-6 secretion and induced insulin resistance in adipocyte: prevention by rosiglitazone. *Biochem Biophys Res Commun* 311:372-379.
 153. Landin K, Lonnroth P, Krotkiewski M, Holm G, Smith U. 1990. Increased insulin resistance and fat cell lipolysis in obese but not lean women with a high waist/hip ratio. *Eur J Clin Invest* 20:530-535.
 154. Langohr IM, HogenEsch H, Stevenson GW, Sturek M. 2008. Vascular-associated lymphoid tissue in swine (*Sus scrofa*). *Comp Med* 58:168-173.
 155. Larsen MO, Juhl CB, Porksen N, Gotfredsen CF, Carr RD, Ribel U, Wilken M, Rolin B. 2005. Beta-cell function and islet morphology in normal, obese, and obese beta-cell mass-reduced Gottingen minipigs. *Am J Physiol Endocrinol Metab* 288:E412-421.
 156. Larsen MO, Rolin B. 2004. Use of the Gottingen minipig as a model of diabetes, with special focus on type 1 diabetes research. *ILAR J* 45:303-313.
 157. Larsen MO, Rolin B, Raun K, Bjerre Knudsen L, Gotfredsen CF, Bock T. 2007. Evaluation of beta-cell mass and function in the Gottingen minipig. *Diabetes Obes Metab* 9 Suppl 2:170-179.
 158. Larsen MO, Rolin B, Wilken M, Carr RD, Svendsen O. 2002. High-fat high-energy feeding impairs fasting glucose and increases fasting insulin levels in the Gottingen minipig: results from a pilot study. *Ann N Y Acad Sci* 967:414-423.
 159. Lassaletta AD, Chu LM, Robich MP, Elmadhun NY, Feng J, Burgess TA, Laham RJ, Sturek M, Sellke FW. 2012. Overfed Ossabaw swine with early stage metabolic syndrome have normal coronary collateral development in response to chronic ischemia. *Basic Res Cardiol* 107:1-11.
 160. Laugerette F, Vors C, Geloën A, Chauvin MA, Soulage C, Lambert-Porcheron S, Peretti N, Alligier M, Burcelin R, Laville M, Vidal H, Michalski MC. 2011. Emulsified lipids increase endotoxemia: possible role in early postprandial low-grade inflammation. *J Nutr Biochem* 22:53-59.
 161. Lee J, Xu Y, Lu L, Bergman B, Leitner JW, Greyson C, Draznin B, Schwartz GG. 2010. Multiple abnormalities of myocardial insulin signaling in a porcine model of diet-induced obesity. *Am J Physiol Heart Circ Physiol* 298:H310-319.

162. Lee JY, Plakidas A, Lee WH, Heikkinen A, Chanmugam P, Bray G, Hwang DH. 2003. Differential modulation of Toll-like receptors by fatty acids: preferential inhibition by n-3 polyunsaturated fatty acids. *J Lipid Res* 44:479-486.
163. Lee JY, Sohn KH, Rhee SH, Hwang D. 2001. Saturated fatty acids, but not unsaturated fatty acids, induce the expression of cyclooxygenase-2 mediated through Toll-like receptor 4. *J Biol Chem* 276:16683-16689.
164. Lee KU, Lee HK, Koh CS, Min HK. 1988. Artificial induction of intravascular lipolysis by lipid-heparin infusion leads to insulin resistance in man. *Diabetologia* 31:285-290.
165. Lee L, Alloosh M, Saxena R, Van Alstine W, Watkins BA, Klaunig JE, Sturek M, Chalasani N. 2009. Nutritional model of steatohepatitis and metabolic syndrome in the Ossabaw miniature swine. *Hepatology* 50:56-67.
166. Li L, Yang G, Li Q, Tang Y, Li K. 2006. High-fat- and lipid-induced insulin resistance in rats: the comparison of glucose metabolism, plasma resistin and adiponectin levels. *Ann Nutr Metab* 50:499-505.
167. Li Q, Yin W, Cai M, Liu Y, Hou H, Shen Q, Zhang C, Xiao J, Hu X, Wu Q, Funaki M, Nakaya Y. 2010. NO-1886 suppresses diet-induced insulin resistance and cholesterol accumulation through STAT5-dependent upregulation of IGF1 and CYP7A1. *J Endocrinol* 204:47-56.
168. Li Z, Woollard JR, Wang S, Korsmo MJ, Ebrahimi B, Grande JP, Textor SC, Lerman A, Lerman LO. 2011. Increased glomerular filtration rate in early metabolic syndrome is associated with renal adiposity and microvascular proliferation. *Am J Physiol Renal Physiol* 301:F1078-1087.
169. Lin YH, Salem N, Jr. 2007. Whole body distribution of deuterated linoleic and alpha-linolenic acids and their metabolites in the rat. *J Lipid Res* 48:2709-2724.
170. Litten-Brown JC, Corson AM, Clarke L. 2010. Porcine models for the metabolic syndrome, digestive and bone disorders: a general overview. *Animal* 4:899-920.
171. Liu Y, Wang Z, Yin W, Li Q, Cai M, Zhang C, Xiao J, Hou H, Li H, Zu X. 2007. Severe insulin resistance and moderate glomerulosclerosis in a minipig model induced by high-fat/ high-sucrose/ high-cholesterol diet. *Exp Anim* 56:11-20.
172. Lloyd PG, Fang MY, Brisbin IL, Andersson L, Sturek M. 2006. AMP kinase gene mutation is consistent with a thrifty phenotype (metabolic syndrome) in a population of feral swine. *Faseb J* 20:A299-A299.
173. Long Q, Lei T, Feng B, Yin C, Jin D, Wu Y, Zhu X, Chen X, Gan L, Yang Z. 2010. Peroxisome proliferator-activated receptor-gamma increases adiponectin secretion via

- transcriptional repression of endoplasmic reticulum chaperone protein ERp44. *Endocrinology* 151:3195-3203.
174. Long SD, Pekala PH. 1996. Lipid mediators of insulin resistance: ceramide signalling down-regulates GLUT4 gene transcription in 3T3-L1 adipocytes. *Biochem J* 319 (Pt 1):179-184.
 175. Long X, Mokolke EA, Neeb ZP, Alloosh M, Edwards JM, Sturek M. 2010. Adenosine receptor regulation of coronary blood flow in Ossabaw miniature swine. *J Pharmacol Exp Ther* 335:781-787.
 176. Loos RJ, Lindgren CM, Li S, Wheeler E, Zhao JH, Prokopenko I, Inouye M, Freathy RM, Attwood AP, Beckmann JS, Berndt SI, Jacobs KB, Chanock SJ, Hayes RB, Bergmann S, Bennett AJ, Bingham SA, Bochud M, Brown M, Cauchi S, Connell JM, Cooper C, Smith GD, Day I, Dina C, De S, Dermitzakis ET, Doney AS, Elliott KS, Elliott P, Evans DM, Sadaf Farooqi I, Froguel P, Ghorri J, Groves CJ, Gwilliam R, Hadley D, Hall AS, Hattersley AT, Hebebrand J, Heid IM, Lamina C, Gieger C, Illig T, Meitinger T, Wichmann HE, Herrera B, Hinney A, Hunt SE, Jarvelin MR, Johnson T, Jolley JD, Karpe F, Keniry A, Khaw KT, Luben RN, Mangino M, Marchini J, McArdle WL, McGinnis R, Meyre D, Munroe PB, Morris AD, Ness AR, Neville MJ, Nica AC, Ong KK, O'Rahilly S, Owen KR, Palmer CN, Papadakis K, Potter S, Pouta A, Qi L, Randall JC, Rayner NW, Ring SM, Sandhu MS, Scherag A, Sims MA, Song K, Soranzo N, Speliotes EK, Syddall HE, Teichmann SA, Timpson NJ, Tobias JH, Uda M, Vogel CI, Wallace C, Waterworth DM, Weedon MN, Willer CJ, Wraight, Yuan X, Zeggini E, Hirschhorn JN, Strachan DP, Ouwehand WH, Caulfield MJ, Samani NJ, Frayling TM, Vollenweider P, Waeber G, Mooser V, Deloukas P, McCarthy MI, Wareham NJ, Barroso I, Kraft P, Hankinson SE, Hunter DJ, Hu FB, Lyon HN, Voight BF, Ridderstrale M, Groop L, Scheet P, Sanna S, Abecasis GR, Albai G, Nagaraja R, Schlessinger D, Jackson AU, Tuomilehto J, Collins FS, Boehnke M, Mohlke KL. 2008. Common variants near MC4R are associated with fat mass, weight and risk of obesity. *Nat Genet* 40:768-775.
 177. Lumeng CN, DelProposto JB, Westcott DJ, Saltiel AR. 2008. Phenotypic switching of adipose tissue macrophages with obesity is generated by spatiotemporal differences in macrophage subtypes. *Diabetes* 57:3239-3246.
 178. MacDougald OA, Burant CF. 2007. The rapidly expanding family of adipokines. *Cell Metab* 6:159-161.
 179. Madsen MB, Birck MM, Fredholm M, Cirera S. 2010. Expression studies of the obesity candidate gene FTO in pig. *Anim Biotechnol* 21:51-63.
 180. Maeda K, Okubo K, Shimomura I, Funahashi T, Matsuzawa Y, Matsubara K. 1996. cDNA cloning and expression of a novel adipose specific collagen-like factor, apM1

- (AdiPose Most abundant Gene transcript 1). *Biochem Biophys Res Commun* 221:286-289.
181. Marchesini G, Brizi M, Morselli-Labate AM, Bianchi G, Bugianesi E, McCullough AJ, Forlani G, Melchionda N. 1999. Association of nonalcoholic fatty liver disease with insulin resistance. *Am J Med* 107:450-455.
 182. Martin RJ, Gobble JL, Hartsock TH, Graves HB, Ziegler JH. 1973. Characterization of an obese syndrome in the pig. *Proc Soc Exp Biol Med* 143:198-203.
 183. Martin RJ, Herbein JH. 1976. A comparison of the enzyme levels and the in vitro utilization of various substrates for lipogenesis in pair-fed lean and obese pigs. *Proc Soc Exp Biol Med* 151:231-235.
 184. Martin RJ, Wangsness PJ, Gahagan JH. 1978. Diurnal changes in serum metabolites and hormones in lean and obese Zucker rats. *Horm Metab Res* 10:187-192.
 185. Martinovic I, Abegunewardene N, Seul M, Vosseler M, Horstick G, Buerke M, Darius H, Lindemann S. 2005. Elevated monocyte chemoattractant protein-1 serum levels in patients at risk for coronary artery disease. *Circ J* 69:1484-1489.
 186. Maury E, Noel L, Detry R, Brichard SM. 2009. In vitro hyperresponsiveness to tumor necrosis factor-alpha contributes to adipokine dysregulation in omental adipocytes of obese subjects. *J Clin Endocrinol Metab* 94:1393-1400.
 187. Mayer JJ, Brisbin IL. 1991. *Wild pigs in the United States : their history, comparative morphology, and current status*. Athens: University of Georgia Press.
 188. McDonald TO, Gerrity RG, Jen C, Chen HJ, Wark K, Wight TN, Chait A, O'Brien KD. 2007. Diabetes and arterial extracellular matrix changes in a porcine model of atherosclerosis. *J Histochem Cytochem* 55:1149-1157.
 189. McQuaid SE, Hodson L, Neville MJ, Dennis AL, Cheeseman J, Humphreys SM, Ruge T, Gilbert M, Fielding BA, Frayn KN, Karpe F. 2011. Downregulation of adipose tissue fatty acid trafficking in obesity: a driver for ectopic fat deposition? *Diabetes* 60:47-55.
 190. Mehta NN, McGillicuddy FC, Anderson PD, Hinkle CC, Shah R, Pruscino L, Tabita-Martinez J, Sellers KF, Rickels MR, Reilly MP. 2010. Experimental endotoxemia induces adipose inflammation and insulin resistance in humans. *Diabetes* 59:172-181.
 191. Meigs JB, Wilson PW, Fox CS, Vasan RS, Nathan DM, Sullivan LM, D'Agostino RB. 2006. Body mass index, metabolic syndrome, and risk of type 2 diabetes or cardiovascular disease. *J Clin Endocrinol Metab* 91:2906-2912.

192. Mestas J, Hughes CC. 2004. Of mice and not men: differences between mouse and human immunology. *J Immunol* 172:2731-2738.
193. Micallef MA, Munro IA, Garg ML. 2009. An inverse relationship between plasma n-3 fatty acids and C-reactive protein in healthy individuals. *Eur J Clin Nutr* 63:1154-1156.
194. Miles PD, Romeo OM, Higo K, Cohen A, Rafaat K, Olefsky JM. 1997. TNF-alpha-induced insulin resistance in vivo and its prevention by troglitazone. *Diabetes* 46:1678-1683.
195. Mohamed-Ali V, Goodrick S, Rawesh A, Katz DR, Miles JM, Yudkin JS, Klein S, Coppack SW. 1997. Subcutaneous adipose tissue releases interleukin-6, but not tumor necrosis factor-alpha, in vivo. *J Clin Endocrinol Metab* 82:4196-4200.
196. Mokolke EA, Hu Q, Song M, Toro L, Reddy HK, Sturek M. 2003. Altered functional coupling of coronary K⁺ channels in diabetic dyslipidemic pigs is prevented by exercise. *J Appl Physiol* 95:1179-1193.
197. Monteiro R, Azevedo I. 2010. Chronic inflammation in obesity and the metabolic syndrome. *Mediators Inflamm* 2010.
198. Morris DL, Singer K, Lumeng CN. 2011. Adipose tissue macrophages: phenotypic plasticity and diversity in lean and obese states. *Curr Opin Clin Nutr Metab Care* 14:341-346.
199. Moshage HJ, Roelofs HM, van Pelt JF, Hazenberg BP, van Leeuwen MA, Limburg PC, Aarden LA, Yap SH. 1988. The effect of interleukin-1, interleukin-6 and its interrelationship on the synthesis of serum amyloid A and C-reactive protein in primary cultures of adult human hepatocytes. *Biochem Biophys Res Commun* 155:112-117.
200. Mottillo EP, Shen XJ, Granneman JG. 2010. beta3-adrenergic receptor induction of adipocyte inflammation requires lipolytic activation of stress kinases p38 and JNK. *Biochim Biophys Acta* 1801:1048-1055.
201. Natarajan R, Gerrity RG, Gu JL, Lanting L, Thomas L, Nadler JL. 2002. Role of 12-lipoxygenase and oxidant stress in hyperglycaemia-induced acceleration of atherosclerosis in a diabetic pig model. *Diabetologia* 45:125-133.
202. Nawrocki AR, Rajala MW, Tomas E, Pajvani UB, Saha AK, Trumbauer ME, Pang Z, Chen AS, Ruderman NB, Chen H, Rossetti L, Scherer PE. 2006. Mice lacking adiponectin show decreased hepatic insulin sensitivity and reduced responsiveness to peroxisome proliferator-activated receptor gamma agonists. *J Biol Chem* 281:2654-2660.

203. Neeb ZP, Edwards JM, Alloosh M, Long X, Mokelke EA, Sturek M. 2010. Metabolic syndrome and coronary artery disease in Ossabaw compared with Yucatan swine. *Comp Med* 60:300-315.
204. Nguyen NT, Nguyen XM, Lane J, Wang P. 2011. Relationship between obesity and diabetes in a US adult population: findings from the National Health and Nutrition Examination Survey, 1999-2006. *Obes Surg* 21:351-355.
205. Novak TE, Babcock TA, Jho DH, Helton WS, Espat NJ. 2003. NF-kappa B inhibition by omega -3 fatty acids modulates LPS-stimulated macrophage TNF-alpha transcription. *Am J Physiol Lung Cell Mol Physiol* 284:L84-89.
206. Olefsky JM, Glass CK. 2010. Macrophages, inflammation, and insulin resistance. *Annu Rev Physiol* 72:219-246.
207. Oster RT, Tishinsky JM, Yuan Z, Robinson LE. 2010. Docosahexaenoic acid increases cellular adiponectin mRNA and secreted adiponectin protein, as well as PPARgamma mRNA, in 3T3-L1 adipocytes. *Appl Physiol Nutr Metab* 35:783-789.
208. Otieno CJ, Bastiaansen J, Ramos AM, Rothschild MF. 2005. Mapping and association studies of diabetes related genes in the pig. *Anim Genet* 36:36-42.
209. Otis CR, Wamhoff BR, Sturek M. 2003. Hyperglycemia-induced insulin resistance in diabetic dyslipidemic Yucatan swine. *Comp Med* 53:53-64.
210. Ouchi N, Kihara S, Funahashi T, Nakamura T, Nishida M, Kumada M, Okamoto Y, Ohashi K, Nagaretani H, Kishida K, Nishizawa H, Maeda N, Kobayashi H, Hiraoka H, Matsuzawa Y. 2003. Reciprocal association of C-reactive protein with adiponectin in blood stream and adipose tissue. *Circulation* 107:671-674.
211. Ouchi N, Walsh K. 2007. Adiponectin as an anti-inflammatory factor. *Clin Chim Acta* 380:24-30.
212. Ovilo C, Fernandez A, Fernandez AI, Folch JM, Varona L, Benitez R, Nunez Y, Rodriguez C, Silio L. 2010. Hypothalamic expression of porcine leptin receptor (LEPR), neuropeptide Y (NPY), and cocaine- and amphetamine-regulated transcript (CART) genes is influenced by LEPR genotype. *Mamm Genome* 21:583-591.
213. Ozes ON, Akca H, Mayo LD, Gustin JA, Maehama T, Dixon JE, Donner DB. 2001. A phosphatidylinositol 3-kinase/Akt/mTOR pathway mediates and PTEN antagonizes tumor necrosis factor inhibition of insulin signaling through insulin receptor substrate-1. *Proc Natl Acad Sci U S A* 98:4640-4645.

214. Pacher P, Beckman JS, Liaudet L. 2007. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* 87:315-424.
215. Pasceri V, Cheng JS, Willerson JT, Yeh ET. 2001. Modulation of C-reactive protein-mediated monocyte chemoattractant protein-1 induction in human endothelial cells by anti-atherosclerosis drugs. *Circulation* 103:2531-2534.
216. Payne GA, Borbouse L, Kumar S, Neeb Z, Alloosh M, Sturek M, Tune JD. 2010. Epicardial perivascular adipose-derived leptin exacerbates coronary endothelial dysfunction in metabolic syndrome via a protein kinase C-beta pathway. *Arterioscler Thromb Vasc Biol* 30:1711-1717.
217. Pepys MB, Hirschfield GM. 2003. C-reactive protein: a critical update. *J Clin Invest* 111:1805-1812.
218. Peraldi P, Xu M, Spiegelman BM. 1997. Thiazolidinediones block tumor necrosis factor-alpha-induced inhibition of insulin signaling. *J Clin Invest* 100:1863-1869.
219. Pessler D, Rudich A, Bashan N. 2001. Oxidative stress impairs nuclear proteins binding to the insulin responsive element in the GLUT4 promoter. *Diabetologia* 44:2156-2164.
220. Phillips DI, Caddy S, Ilic V, Fielding BA, Frayn KN, Borthwick AC, Taylor R. 1996. Intramuscular triglyceride and muscle insulin sensitivity: evidence for a relationship in nondiabetic subjects. *Metabolism* 45:947-950.
221. Ponnampalam EN, Lewandowski P, Nesaratnam K, Dunshea FR, Gill H. 2011. Differential effects of natural palm oil, chemically- and enzymatically-modified palm oil on weight gain, blood lipid metabolites and fat deposition in a pediatric pig model. *Nutr J* 10:53.
222. Prins JB, Niesler CU, Winterford CM, Bright NA, Siddle K, O'Rahilly S, Walker NI, Cameron DP. 1997. Tumor necrosis factor-alpha induces apoptosis of human adipose cells. *Diabetes* 46:1939-1944.
223. Punyadeera C, van der Merwe MT, Crowther NJ, Toman M, Immelman AR, Schlaphoff GP, Gray IP. 2001. Weight-related differences in glucose metabolism and free fatty acid production in two South African population groups. *Int J Obes Relat Metab Disord* 25:1196-1205.
224. Pussinen PJ, Havulinna AS, Lehto M, Sundvall J, Salomaa V. 2011. Endotoxemia is associated with an increased risk of incident diabetes. *Diabetes Care* 34:392-397.
225. Raetz CR. 1990. Biochemistry of endotoxins. *Annu Rev Biochem* 59:129-170.

226. Raun K, von Voss P, Knudsen LB. 2007. Liraglutide, a once-daily human glucagon-like peptide-1 analog, minimizes food intake in severely obese minipigs. *Obesity (Silver Spring)* 15:1710-1716.
227. Reinehr T, Kiess W, Andler W. 2005. Insulin sensitivity indices of glucose and free fatty acid metabolism in obese children and adolescents in relation to serum lipids. *Metabolism* 54:397-402.
228. Robich MP, Chu LM, Chaudray M, Nezafat R, Han Y, Clements RT, Laham RJ, Manning WJ, Coady MA, Sellke FW. 2010. Anti-angiogenic effect of high-dose resveratrol in a swine model of metabolic syndrome. *Surgery* 148:453-462.
229. Robich MP, Osipov RM, Chu LM, Han Y, Feng J, Nezafat R, Clements RT, Manning WJ, Sellke FW. 2011. Resveratrol modifies risk factors for coronary artery disease in swine with metabolic syndrome and myocardial ischemia. *Eur J Pharmacol* 664:45-53.
230. Robinson C, Tamborlane WV, Maggs DG, Enoksson S, Sherwin RS, Silver D, Shulman GI, Caprio S. 1998. Effect of insulin on glycerol production in obese adolescents. *Am J Physiol* 274:E737-743.
231. Roshick C, Wood H, Caldwell HD, McClarty G. 2006. Comparison of gamma interferon-mediated antichlamydial defense mechanisms in human and mouse cells. *Infect Immun* 74:225-238.
232. Rotter V, Nagaev I, Smith U. 2003. Interleukin-6 (IL-6) induces insulin resistance in 3T3-L1 adipocytes and is, like IL-8 and tumor necrosis factor-alpha, overexpressed in human fat cells from insulin-resistant subjects. *J Biol Chem* 278:45777-45784.
233. Ruan H, Hacohen N, Golub TR, Van Parijs L, Lodish HF. 2002. Tumor necrosis factor-alpha suppresses adipocyte-specific genes and activates expression of preadipocyte genes in 3T3-L1 adipocytes: nuclear factor-kappaB activation by TNF-alpha is obligatory. *Diabetes* 51:1319-1336.
234. Ruan H, Pownall HJ, Lodish HF. 2003. Troglitazone antagonizes tumor necrosis factor-alpha-induced reprogramming of adipocyte gene expression by inhibiting the transcriptional regulatory functions of NF-kappaB. *J Biol Chem* 278:28181-28192.
235. Rudich A, Tirosh A, Potashnik R, Hemi R, Kanety H, Bashan N. 1998. Prolonged oxidative stress impairs insulin-induced GLUT4 translocation in 3T3-L1 adipocytes. *Diabetes* 47:1562-1569.
236. Rui L, Aguirre V, Kim JK, Shulman GI, Lee A, Corbould A, Dunaif A, White MF. 2001. Insulin/IGF-1 and TNF-alpha stimulate phosphorylation of IRS-1 at inhibitory Ser307 via distinct pathways. *J Clin Invest* 107:181-189.

237. Ryan JG. 2009. Cost and policy implications from the increasing prevalence of obesity and diabetes mellitus. *Gend Med* 6 Suppl 1:86-108.
238. Sabin MA, Yau SW, Russo VC, Clarke IJ, Dunshea FR, Chau J, Cox M, Werther GA. 2011. Dietary monounsaturated fat in early life regulates IGFBP2: implications for fat mass accretion and insulin sensitivity. *Obesity (Silver Spring)* 19:2374-2381.
239. Scherer PE, Williams S, Fogliano M, Baldini G, Lodish HF. 1995. A novel serum protein similar to C1q, produced exclusively in adipocytes. *J Biol Chem* 270:26746-26749.
240. Schwartz EA, Zhang WY, Karnik SK, Borwege S, Anand VR, Laine PS, Su Y, Reaven PD. 2010. Nutrient modification of the innate immune response: a novel mechanism by which saturated fatty acids greatly amplify monocyte inflammation. *Arterioscler Thromb Vasc Biol* 30:802-808.
241. Sebert SP, Lecannu G, Kozlowski F, Siliart B, Bard JM, Krempf M, Champ MM. 2005. Childhood obesity and insulin resistance in a Yucatan mini-piglet model: putative roles of IGF-1 and muscle PPARs in adipose tissue activity and development. *Int J Obes (Lond)* 29:324-333.
242. Shamseddeen H, Getty JZ, Hamdallah IN, Ali MR. 2011. Epidemiology and economic impact of obesity and type 2 diabetes. *Surg Clin North Am* 91:1163-1172, vii.
243. Shapiro H, Lutaty A, Ariel A. 2011. Macrophages, meta-inflammation, and immuno-metabolism. *ScientificWorldJournal* 11:2509-2529.
244. Shen WH, Sturek M, Liechty EA. 2007. Molecular mechanisms for insulin resistance in Ossabaw miniature swine. *Faseb J* 21:A1053-A1053.
245. Shi H, Kokoeva MV, Inouye K, Tzameli I, Yin H, Flier JS. 2006. TLR4 links innate immunity and fatty acid-induced insulin resistance. *J Clin Invest* 116:3015-3025.
246. Shimabukuro M, Zhou YT, Levi M, Unger RH. 1998. Fatty acid-induced beta cell apoptosis: a link between obesity and diabetes. *Proc Natl Acad Sci U S A* 95:2498-2502.
247. Shin MJ, Hyun YJ, Kim OY, Kim JY, Jang Y, Lee JH. 2006. Weight loss effect on inflammation and LDL oxidation in metabolically healthy but obese (MHO) individuals: low inflammation and LDL oxidation in MHO women. *Int J Obes (Lond)* 30:1529-1534.

248. Sims EA. 2001. Are there persons who are obese, but metabolically healthy? *Metabolism* 50:1499-1504.
249. Skoog T, Dichtl W, Boquist S, Skoglund-Andersson C, Karpe F, Tang R, Bond MG, de Faire U, Nilsson J, Eriksson P, Hamsten A. 2002. Plasma tumour necrosis factor-alpha and early carotid atherosclerosis in healthy middle-aged men. *Eur Heart J* 23:376-383.
250. Skurk T, Alberti-Huber C, Herder C, Hauner H. 2007. Relationship between adipocyte size and adipokine expression and secretion. *J Clin Endocrinol Metab* 92:1023-1033.
251. Smith WL. 2005. Cyclooxygenases, peroxide tone and the allure of fish oil. *Curr Opin Cell Biol* 17:174-182.
252. Song MJ, Kim KH, Yoon JM, Kim JB. 2006. Activation of Toll-like receptor 4 is associated with insulin resistance in adipocytes. *Biochem Biophys Res Commun* 346:739-745.
253. Spurlock ME, Gabler NK. 2008. The development of porcine models of obesity and the metabolic syndrome. *J Nutr* 138:397-402.
254. Stefan N, Kantartzis K, Machann J, Schick F, Thamer C, Rittig K, Balletshofer B, Machicao F, Fritsche A, Haring HU. 2008. Identification and characterization of metabolically benign obesity in humans. *Arch Intern Med* 168:1609-1616.
255. Stephens JM, Lee J, Pilch PF. 1997. Tumor necrosis factor-alpha-induced insulin resistance in 3T3-L1 adipocytes is accompanied by a loss of insulin receptor substrate-1 and GLUT4 expression without a loss of insulin receptor-mediated signal transduction. *J Biol Chem* 272:971-976.
256. Stillwell W, Wassall SR. 2003. Docosaehaenoic acid: membrane properties of a unique fatty acid. *Chem Phys Lipids* 126:1-27.
257. Straczkowski M, Dzienis-Straczkowska S, Stepien A, Kowalska I, Szelachowska M, Kinalska I. 2002. Plasma interleukin-8 concentrations are increased in obese subjects and related to fat mass and tumor necrosis factor-alpha system. *J Clin Endocrinol Metab* 87:4602-4606.
258. Stribling HL, Brisbin IL, Jr., Sweeney JR, Stribling LA. 1984. Body Fat Reserves and Their Prediction in Two Populations of Feral Swine. *The Journal of Wildlife Management* 48:635-639.

259. Strissel KJ, Stancheva Z, Miyoshi H, Perfield JW, 2nd, DeFuria J, Jick Z, Greenberg AS, Obin MS. 2007. Adipocyte death, adipose tissue remodeling, and obesity complications. *Diabetes* 56:2910-2918.
260. Suganami T, Tanimoto-Koyama K, Nishida J, Itoh M, Yuan X, Mizuarai S, Kotani H, Yamaoka S, Miyake K, Aoe S, Kamei Y, Ogawa Y. 2007. Role of the Toll-like receptor 4/NF-kappaB pathway in saturated fatty acid-induced inflammatory changes in the interaction between adipocytes and macrophages. *Arterioscler Thromb Vasc Biol* 27:84-91.
261. Suzuki LA, Poot M, Gerrity RG, Bornfeldt KE. 2001. Diabetes accelerates smooth muscle accumulation in lesions of atherosclerosis: lack of direct growth-promoting effects of high glucose levels. *Diabetes* 50:851-860.
262. Swindle MM. 2007. *Swine in the laboratory : surgery, anesthesia, imaging, and experimental techniques*. Boca Raton: CRC Press.
263. Switonski M, Stachowiak M, Cieslak J, Bartz M, Grzes M. 2010. Genetics of fat tissue accumulation in pigs: a comparative approach. *J Appl Genet* 51:153-168.
264. Szalai AJ, McCrory MA. 2002. Varied biologic functions of C-reactive protein: lessons learned from transgenic mice. *Immunol Res* 26:279-287.
265. Tacke F, Randolph GJ. 2006. Migratory fate and differentiation of blood monocyte subsets. *Immunobiology* 211:609-618.
266. Taha C, Klip A. 1999. The insulin signaling pathway. *J Membr Biol* 169:1-12.
267. Tateya S, Tamori Y, Kawaguchi T, Kanda H, Kasuga M. 2010. An increase in the circulating concentration of monocyte chemoattractant protein-1 elicits systemic insulin resistance irrespective of adipose tissue inflammation in mice. *Endocrinology* 151:971-979.
268. Temelkova-Kurktschiev T, Stefanov T. 2012. Lifestyle and genetics in obesity and type 2 diabetes. *Exp Clin Endocrinol Diabetes* 120:1-6.
269. Teruel T, Hernandez R, Lorenzo M. 2001. Ceramide mediates insulin resistance by tumor necrosis factor-alpha in brown adipocytes by maintaining Akt in an inactive dephosphorylated state. *Diabetes* 50:2563-2571.
270. Thoma-Uszynski S, Stenger S, Takeuchi O, Ochoa MT, Engele M, Sieling PA, Barnes PF, Rollinghoff M, Bolcskei PL, Wagner M, Akira S, Norgard MV, Belisle JT, Godowski PJ, Bloom BR, Modlin RL. 2001. Induction of direct antimicrobial activity through mammalian toll-like receptors. *Science* 291:1544-1547.

271. Todoric J, Loffler M, Huber J, Bilban M, Reimers M, Kadl A, Zeyda M, Waldhausl W, Stulnig TM. 2006. Adipose tissue inflammation induced by high-fat diet in obese diabetic mice is prevented by n-3 polyunsaturated fatty acids. *Diabetologia* 49:2109-2119.
272. Tomas E, Tsao TS, Saha AK, Murrey HE, Zhang Cc C, Itani SI, Lodish HF, Ruderman NB. 2002. Enhanced muscle fat oxidation and glucose transport by ACRP30 globular domain: acetyl-CoA carboxylase inhibition and AMP-activated protein kinase activation. *Proc Natl Acad Sci U S A* 99:16309-16313.
273. Tsukumo DM, Carvalho-Filho MA, Carvalheira JB, Prada PO, Hirabara SM, Schenka AA, Araujo EP, Vassallo J, Curi R, Velloso LA, Saad MJ. 2007. Loss-of-function mutation in Toll-like receptor 4 prevents diet-induced obesity and insulin resistance. *Diabetes* 56:1986-1998.
274. Ueki K, Kondo T, Kahn CR. 2004. Suppressor of cytokine signaling 1 (SOCS-1) and SOCS-3 cause insulin resistance through inhibition of tyrosine phosphorylation of insulin receptor substrate proteins by discrete mechanisms. *Mol Cell Biol* 24:5434-5446.
275. Val-Laillet D, Guerin S, Malbert CH. 2010. Slower eating rate is independent to gastric emptying in obese minipigs. *Physiol Behav* 101:462-468.
276. van Hall G, Steensberg A, Sacchetti M, Fischer C, Keller C, Schjerling P, Hiscock N, Moller K, Saltin B, Febbraio MA, Pedersen BK. 2003. Interleukin-6 stimulates lipolysis and fat oxidation in humans. *J Clin Endocrinol Metab* 88:3005-3010.
277. Vozarova B, Weyer C, Hanson K, Tataranni PA, Bogardus C, Pratley RE. 2001. Circulating interleukin-6 in relation to adiposity, insulin action, and insulin secretion. *Obes Res* 9:414-417.
278. Wang B, Trayhurn P. 2006. Acute and prolonged effects of TNF-alpha on the expression and secretion of inflammation-related adipokines by human adipocytes differentiated in culture. *Pflugers Arch* 452:418-427.
279. Wang YC, McPherson K, Marsh T, Gortmaker SL, Brown M. 2011. Health and economic burden of the projected obesity trends in the USA and the UK. *Lancet* 378:815-825.
280. Wang ZV, Schraw TD, Kim JY, Khan T, Rajala MW, Follenzi A, Scherer PE. 2007. Secretion of the adipocyte-specific secretory protein adiponectin critically depends on thiol-mediated protein retention. *Mol Cell Biol* 27:3716-3731.
281. Wangsness PJ, Gobble JL, Sherrett GW. 1980. Feeding-Behavior of Lean and Obese Pigs. *Physiol Behav* 24:407-410.

282. Wangsness PJ, Martin RJ, Gahagan JH. 1977. Insulin and growth hormone in lean and obese pigs. *Am J Physiol* 233:E104-108.
283. Wangsness PJ, Martin RJ, Gatchel BB. 1980. Insulin induced growth hormone response in fast-growing, lean and in slow-growing, obese pigs. *Growth* 44:318-326.
284. Wassall SR, Stillwell W. 2008. Docosaehaenoic acid domains: the ultimate non-raft membrane domain. *Chem Phys Lipids* 153:57-63.
285. Weigert C, Hennige AM, Lehmann R, Brodbeck K, Baumgartner F, Schauble M, Haring HU, Schleicher ED. 2006. Direct cross-talk of interleukin-6 and insulin signal transduction via insulin receptor substrate-1 in skeletal muscle cells. *J Biol Chem* 281:7060-7067.
286. Weisberg SP, Hunter D, Huber R, Lemieux J, Slaymaker S, Vaddi K, Charo I, Leibel RL, Ferrante AW, Jr. 2006. CCR2 modulates inflammatory and metabolic effects of high-fat feeding. *J Clin Invest* 116:115-124.
287. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW, Jr. 2003. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* 112:1796-1808.
288. Weiss R, Taksali SE, Dufour S, Yeckel CW, Papademetris X, Cline G, Tamborlane WV, Dziura J, Shulman GI, Caprio S. 2005. The "obese insulin-sensitive" adolescent: importance of adiponectin and lipid partitioning. *J Clin Endocrinol Metab* 90:3731-3737.
289. Wentworth JM, Naselli G, Brown WA, Doyle L, Phipson B, Smyth GK, Wabitsch M, O'Brien PE, Harrison LC. 2010. Pro-inflammatory CD11c+CD206+ adipose tissue macrophages are associated with insulin resistance in human obesity. *Diabetes* 59:1648-1656.
290. Witczak CA, Mokolke EA, Boullion R, Wenzel J, Keisler DH, Sturek M. 2005. Noninvasive measures of body fat percentage in male Yucatan swine. *Comp Med* 55:445-451.
291. Wu J, Wang B, Zhang H, Yu T, Yang G. 2007. Different transcription profiles of SOCS-3, ob and IGF-I genes and their possible correlations in obese and lean pigs. *Acta Biochim Biophys Sin (Shanghai)* 39:305-310.
292. Wu X, Motoshima H, Mahadev K, Stalker TJ, Scalia R, Goldstein BJ. 2003. Involvement of AMP-activated protein kinase in glucose uptake stimulated by the globular domain of adiponectin in primary rat adipocytes. *Diabetes* 52:1355-1363.

293. Wulster-Radcliffe MC, Ajuwon KM, Wang J, Christian JA, Spurlock ME. 2004. Adiponectin differentially regulates cytokines in porcine macrophages. *Biochem Biophys Res Commun* 316:924-929.
294. Xi S, Yin W, Wang Z, Kusunoki M, Lian X, Koike T, Fan J, Zhang Q. 2004. A minipig model of high-fat/high-sucrose diet-induced diabetes and atherosclerosis. *Int J Exp Pathol* 85:223-231.
295. Xu A, Wang Y, Keshaw H, Xu LY, Lam KS, Cooper GJ. 2003. The fat-derived hormone adiponectin alleviates alcoholic and nonalcoholic fatty liver diseases in mice. *J Clin Invest* 112:91-100.
296. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS, Tartaglia LA, Chen H. 2003. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest* 112:1821-1830.
297. Yamaguchi N, Argueta JG, Masuhiro Y, Kagishita M, Nonaka K, Saito T, Hanazawa S, Yamashita Y. 2005. Adiponectin inhibits Toll-like receptor family-induced signaling. *FEBS Lett* 579:6821-6826.
298. Yamauchi T, Hara K, Kubota N, Terauchi Y, Tobe K, Froguel P, Nagai R, Kadowaki T. 2003. Dual roles of adiponectin/Acrp30 in vivo as an anti-diabetic and anti-atherogenic adipokine. *Curr Drug Targets Immune Endocr Metabol Disord* 3:243-254.
299. Yamauchi T, Kamon J, Ito Y, Tsuchida A, Yokomizo T, Kita S, Sugiyama T, Miyagishi M, Hara K, Tsunoda M, Murakami K, Ohteki T, Uchida S, Takekawa S, Waki H, Tsuno NH, Shibata Y, Terauchi Y, Froguel P, Tobe K, Koyasu S, Taira K, Kitamura T, Shimizu T, Nagai R, Kadowaki T. 2003. Cloning of adiponectin receptors that mediate antidiabetic metabolic effects. *Nature* 423:762-769.
300. Yamauchi T, Kamon J, Minokoshi Y, Ito Y, Waki H, Uchida S, Yamashita S, Noda M, Kita S, Ueki K, Eto K, Akanuma Y, Froguel P, Foufelle F, Ferre P, Carling D, Kimura S, Nagai R, Kahn BB, Kadowaki T. 2002. Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med* 8:1288-1295.
301. Yamauchi T, Nio Y, Maki T, Kobayashi M, Takazawa T, Iwabu M, Okada-Iwabu M, Kawamoto S, Kubota N, Kubota T, Ito Y, Kamon J, Tsuchida A, Kumagai K, Kozono H, Hada Y, Ogata H, Tokuyama K, Tsunoda M, Ide T, Murakami K, Awazawa M, Takamoto I, Froguel P, Hara K, Tobe K, Nagai R, Ueki K, Kadowaki T. 2007. Targeted disruption of AdipoR1 and AdipoR2 causes abrogation of adiponectin binding and metabolic actions. *Nat Med* 13:332-339.

302. Yang Y, Ju D, Zhang M, Yang G. 2008. Interleukin-6 stimulates lipolysis in porcine adipocytes. *Endocrine* 33:261-269.
303. Yang Y, Yang G. 2010. Rosiglitazone regulates IL-6-stimulated lipolysis in porcine adipocytes. *Biochem Cell Biol* 88:853-860.
304. Yin W, Liao D, Kusunoki M, Xi S, Tsutsumi K, Wang Z, Lian X, Koike T, Fan J, Yang Y, Tang C. 2004. NO-1886 decreases ectopic lipid deposition and protects pancreatic beta cells in diet-induced diabetic swine. *J Endocrinol* 180:399-408.
305. Yokota T, Oritani K, Takahashi I, Ishikawa J, Matsuyama A, Ouchi N, Kihara S, Funahashi T, Tenner AJ, Tomiyama Y, Matsuzawa Y. 2000. Adiponectin, a new member of the family of soluble defense collagens, negatively regulates the growth of myelomonocytic progenitors and the functions of macrophages. *Blood* 96:1723-1732.
306. Yoon MJ, Lee GY, Chung JJ, Ahn YH, Hong SH, Kim JB. 2006. Adiponectin increases fatty acid oxidation in skeletal muscle cells by sequential activation of AMP-activated protein kinase, p38 mitogen-activated protein kinase, and peroxisome proliferator-activated receptor alpha. *Diabetes* 55:2562-2570.
307. Yudkin JS, Juhan-Vague I, Hawe E, Humphries SE, di Minno G, Margaglione M, Tremoli E, Kooistra T, Morange PE, Lundman P, Mohamed-Ali V, Hamsten A. 2004. Low-grade inflammation may play a role in the etiology of the metabolic syndrome in patients with coronary heart disease: the HIFMECH study. *Metabolism* 53:852-857.
308. Yudkin JS, Stehouwer CD, Emeis JJ, Coppack SW. 1999. C-reactive protein in healthy subjects: associations with obesity, insulin resistance, and endothelial dysfunction: a potential role for cytokines originating from adipose tissue? *Arterioscler Thromb Vasc Biol* 19:972-978.
309. Zangar RC, Davydov DR, Verma S. 2004. Mechanisms that regulate production of reactive oxygen species by cytochrome P450. *Toxicol Appl Pharmacol* 199:316-331.
310. Zeyda M, Farmer D, Todoric J, Aszmann O, Speiser M, Gyori G, Zlabinger GJ, Stulnig TM. 2007. Human adipose tissue macrophages are of an anti-inflammatory phenotype but capable of excessive pro-inflammatory mediator production. *Int J Obes (Lond)* 31:1420-1428.
311. Zeyda M, Saemann MD, Stuhlmeier KM, Mascher DG, Nowotny PN, Zlabinger GJ, Waldhausl W, Stulnig TM. 2005. Polyunsaturated fatty acids block dendritic cell activation and function independently of NF-kappaB activation. *J Biol Chem* 280:14293-14301.
312. Zhang DX, Gutterman DD. 2007. Mitochondrial reactive oxygen species-mediated signaling in endothelial cells. *Am J Physiol Heart Circ Physiol* 292:H2023-2031.

313. Zhang P, Smith R, Chapkin RS, McMurray DN. 2005. Dietary (n-3) polyunsaturated fatty acids modulate murine Th1/Th2 balance toward the Th2 pole by suppression of Th1 development. *J Nutr* 135:1745-1751.
314. Zhao L, Kwon MJ, Huang S, Lee JY, Fukase K, Inohara N, Hwang DH. 2007. Differential modulation of Nods signaling pathways by fatty acids in human colonic epithelial HCT116 cells. *J Biol Chem* 282:11618-11628.
315. Zhu P, Lu L, Xu Y, Schwartz GG. 2000. Troglitazone improves recovery of left ventricular function after regional ischemia in pigs. *Circulation* 101:1165-1171.

Tables

Table 1. *Swine subcutaneous adipose tissue macrophage phenotype^a.*

Marker	Mean (%)	SEM
CD203a+	26.94	2.29
CD163 ⁺	69.82	1.12
CD14 ⁺ CD16 ⁺	67.64	2.60
CD14 ⁺ CD16 ⁻	20.17	2.37
CD14 ⁻ CD16 ⁺	7.52	0.25
CD14 ⁻ CD16 ⁻	4.66	0.38
CD163 ⁻	30.18	1.12
CD14 ⁺ CD16 ⁺	41.15	1.41
CD14 ⁺ CD16 ⁻	21.99	0.81
CD14 ⁻ CD16 ⁺	12.32	0.41
CD14 ⁻ CD16 ⁻	24.54	1.15

^aAdapted from Faris *et al.* (84).

CHAPTER 3: INFLAMMATION AND RESPONSE TO N-3 FATTY ACIDS IN A PORCINE OBESITY MODEL

Accepted by *Comparative Medicine*

Richard J. Faris, Rebecca L. Boddicker, Jennifer Walker-Daniels, Jenny Li, Douglas E. Jones, and Michael E. Spurlock

Abstract

Fatty acids have distinct cellular effects relating to inflammation and insulin sensitivity. Dietary saturated fat activates toll-like receptor-4 which can lead to chronic inflammation, insulin resistance, and adipose tissue macrophage infiltration. Conversely, n-3 fatty acids are generally anti-inflammatory and promote insulin sensitivity, in part via through peroxisome proliferator-activated receptor- γ . Because Ossabaw swine are an attractive biomedical model, we fed Ossabaw pigs either a low fat control (LFC), high fat palm oil (HFP), or high fat palm oil with n-3 fatty acids (HFPn3) diet for 30 weeks to investigate the effect of saturated fats and n-3 fatty acids on obesity-linked inflammatory markers. There was no diet-related induction of the inflammatory markers C-reactive protein, tumor necrosis factor- α , interleukin-6 or interleukin-12. The increase in inflammatory adipose tissue CD16⁺CD14⁺ macrophages by high palm oil was attenuated by n-3 fatty acids. Both high fat diets induced hyperglycemia, without hyperinsulinemia. Insulin sensitivity was decreased in the HFP group, but not the HFPn3 group during an insulin challenge. This effect was not mediated by decreased phosphorylation of PKB. Thus, in a porcine model of obesity, n-3 fatty acids partially attenuate insulin sensitivity, but only marginally change inflammatory status and macrophage phenotype in adipose tissue.

Introduction

Obesity is accompanied by chronic inflammation in adipose tissue, increased circulating concentrations of tumor necrosis factor- α (TNF α), interleukin-6 (IL-6), and C-reactive protein (CRP), and decreased concentrations of adiponectin (2). This chronic inflammation links obesity and the development of insulin resistance (39). Dietary saturated fatty acids promote obesity in part through the induction of inflammation via activation of toll-like receptor-4 (i.e., tlr4, the innate immune receptor for lipopolysaccharide) (28). Absence of functional tlr4 in mice reduces circulating proinflammatory cytokine concentrations and decreases macrophage infiltration into adipose tissue in high fat diet-induced obesity (8; 28; 32). Furthermore, in 3T3 L1 mouse adipocytes, palmitate activates nuclear factor κ B (NF κ B), protein kinase C, and mitogen-activated protein kinase, all of which increase inflammatory cytokine production (1).

For individuals consuming a diet high in saturated fat, a major determinant of health is the n-6:n-3 (fatty acids, omega-6:omega-3) (5-6). Unlike saturated fatty acids, the n-3 polyunsaturated fatty acids (PUFA) eicosapentanoic acid (EPA) and docosahexanoic acid (DHA) exert predominantly anti-inflammatory effects, as is evident in that DHA antagonizes NF κ B activation by palmitate in 3T3 L1 adipocytes (1). In mice, EPA prevents or reverses hyperinsulinemia, hyperglycemia, and increased circulating monocyte chemoattractant protein-1 (MCP-1) (16), and also decreases infiltration of adipose tissue with macrophages (30). Moreover, n-3 PUFA alleviate the decline in serum adiponectin associated with obesity (12; 15; 30), and EPA decreases serum CRP in diabetic patients (26).

Physiological differences between rodents and humans underscore the need for comparative models in biomedical research, and the pig is emerging rapidly as an attractive

model for energy metabolism and obesity. Like humans, pigs are natural omnivores, rely on apo B-100 to shuttle cholesterol in the LDL fraction, and have minimal brown fat retention postnatally. Furthermore, adipose depots in pigs are of sufficient size that multiple assays can be carried out on adipocytes or stromal vascular cells without pooling across depots or animals. Although Ossabaw swine have been used as a model for metabolic syndrome, cardiovascular disease, coronary artery disease, and steatohepatitis (11; 20; 24), very little is known about adipose inflammation in this model. Consequently, we sought to characterize obesity-linked inflammatory markers in the adipose tissue of this novel model, and to test the hypothesis that adding n-3 PUFA to a diet high in saturated fat will attenuate chronic inflammation, protect against diet-induced insulin resistance, and alter phenotypical changes in adipose tissue macrophages.

Methods

Animals and Diets. All animal procedures were conducted at Iowa State University and were approved by the Iowa State University Institutional Animal Care and Use Committee. Two groups (27 total) of mature male and female Ossabaw swine (*Sus scrofa*) (initial body weight 47.4 ± 0.3 kg) were housed in individual pens in a temperature controlled environment with a 12 hour light cycle. Swine were obtained from the Comparative Medicine Program at the Indiana University School of Medicine and Purdue University from a herd that tested negative for *Brucella* spp. and psuedorabies, and showed no clinical evidence of communicable disease upon arrival or during the study. Animals were assigned to one of three dietary treatments: a low fat control diet (LFC, n=9; 3 male and 6 female), a high fat palm oil diet (HFP, n=9; 2 male and 7 female), and a high fat palm oil diet with n-3 supplementation (HFPn3, n=9; 2 male and 7 female) (**Table 2**). Three pigs in

the HFPn3 dietary treatment died during the study. Autopsy results indicated that the pigs did not have any obvious signs of disease. It was noted that severe congestion in the lungs would support a heart condition that was acute and fatal. However, the exact cause of death was unknown. The HFP and HFPn3 diets were fed *ad libitum* and the LFC diet was restricted to maintain a mature lean body weight. The “healthy” control diet (**Table 2**) was designed to limit caloric intake and provide adequate n3 fatty acids through modest inclusions of fish meal, canola oil, and n3 fatty acid supplement (GromegaTM 365). Both high fat diets were high in saturated fatty acids and formulated to be isocaloric. Canola oil and the n3 fatty acid supplement was used to reduce the n6:n3 to be comparable with the control diet. Body weight and feed intake were measured weekly. To confirm the fatty acid profile of each diet, lipids were extracted by the method of Lepage and Roy (21). Fatty acid methyl esters were analyzed by gas chromatography (model 6890; Hewlett-Packard, Palo Alto, CA) fitted with an Omegawax 320 (30 m x 0.32 mm internal diameter, 0.25 μ m) capillary column (Sigma-Aldrich, St. Louis, MO). The injector and detector temperatures were 250°C, and the oven temperature was 200°C.

Insulin Challenge and Tissue Collection. After 30 weeks of dietary treatment, animals were fasted for 12 hours overnight and were anesthetized with telazol:ketamine:xylazine (1:1:1). Thereafter, blood pressure and mean arterial pressure were taken with an oxillometric blood pressure monitor (Cardell, CAS Medical Systems, Inc., Branford, CT) with a 9 cm cuff on the metatarsus. Each measurement was taken three times and averaged. Blood was collected via jugular venipuncture and then animals were challenged with 0.25 IU/kg of porcine insulin (MP Biomedicals, Solon, CA) intravenously as adapted from Xi et al. (37). Glucose measurements were taken from blood obtained by ear

pricks at 0, 5, 10, and 15 minutes post insulin challenge using a glucometer (LifeScan, Milpitas, CA). Pigs were then killed by exsanguination. Subcutaneous (SQ) adipose tissue from back fat above the 10th rib, visceral adipose tissue (VIS) from the ventral intra-abdominal cavity, omental adipose tissue (OM), liver, and longissimus dorsi (LD) from above the 10th rib were collected. Liver and LD were snap frozen in liquid nitrogen. Adipose tissue was divided two ways: snap frozen in liquid nitrogen or stored in ice cold PBS with 0.2% BSA for transport to the laboratory and the isolation of stromal vascular cells (SVC).

SVC were isolated from adipose tissue as follows: adipose tissue was minced finely with a razor blade and digested in Hank's Buffered Salt Solution (HBSS) with 20 mM HEPES (pH 7.4), 3% BSA, and 2 mg/mL Collagenase Type I (Worthington-Biochemical Corporation, Lakewood, NJ) for 45 minutes, 125 rpm, at 37°C. Samples were then centrifuged for 5 min at 1000xg and the supernatant was discarded. The pellet was dissolved in 10 mL of RBC lysis buffer (0.154 M NH₄Cl, 10 mM KHCO₃, 0.1 mM EDTA, pH 7.3) for 5 min to lyse red blood cells, and then 10 mL of PBS was added and samples were passed through a 100 µm filter. The filtrate was centrifuged for 5 min at 1000xg and the pellet was resuspended in HBSS with 20 mM HEPES (pH 7.4).

Flow Cytometry. Up to 1×10^6 SVC were washed in 2 mL of fluorescence-activated cell sorting buffer (FACS, 0.1% sodium azide, 0.1% BSA in PBS). Cells were treated with 10 µL of pig serum for 15 min at 4°C to prevent nonspecific binding. Cells were then incubated with 10 µL of anti-CD203a IgG₁ (1:20) (Mybiosource, LLC, San Diego, CA) or isotype control (mouse IgG₁, κ, 1:20) (eBioscience, Inc., San Diego, CA) for 15 minutes at 4°C and then washed in 2 mL of FACS buffer. Cells were next incubated in 10 µL of goat-

anti mouse IgG₁-FITC (1:50) (SouthernBiotech, Birmingham, AL) for 15 min at 4°C and then washed in 2 mL FACS buffer. Cells were incubated in 10 µL of biotinylated anti-CD16 (1:50) (BD Pharmingen, San Diego, CA) or isotype control (biotinylated mouse IgG₁, κ, 1:50) (BD Pharmingen) for 15 min at 4°C and then washed in 2 mL of FACS buffer. Cells were then incubated in 10 µL of Streptavidin-Peridinin Chlorophyll-a Protein-Cy5.5 (1:50) (BD Pharmingen) and 10 µL of anti-CD14-PE (1:20) (Antigenix America, Huntington Station, NY) or isotype control (PE mouse IgG_{2b}, κ, 1:20) (BD Pharmingen) for 15 minutes at 4°C and then washed in 2 mL of FACS buffer. Cells were then fixed in 200 µL of 1% paraformaldehyde and stored at 4°C until analysis. Analysis was performed on a BD FACScanto flow cytometer (Becton Dickinson, San Jose, CA), and data was analyzed using FlowJo software V8.5.2 (Tree Star, Inc., Ashland, OR).

Serum and Tissue Analyses. Cholesterol, triglycerides, and non-esterified fatty acids (NEFA) were analyzed using Chol Slides (Ortho Clinical Diagnostics, Rochester, NY), Trig Slides (Ortho Clinical Diagnostics), and NEFA-HR (Wako Diagnostics, Richmond, VA), respectively, on a Vitros 5.1 Chemistry Analyzer (Ortho Clinical Diagnostics). Serum insulin and CRP concentrations were measured by EIA and ELISA, respectively (ALPCO, Salem, NH). Serum TNFα, IL-6, IL-10 and IL-12p40 concentrations were measured by ELISA (R & D Systems, Minneapolis, MN). The HOMA-IR was calculated using fasting serum insulin (µU/mL) x fasting blood glucose (mg/dL) divided by 405 (23). Plasma homocysteine concentrations were determined by HPLC as described by Williams et al. (36). Total liver and LD lipid content was determined by the method of Folch et al. (13).

Whole frozen tissue samples were pulverized and homogenized in ice cold buffer (50 mM HEPES pH 7.4, 150 mM NaCl, 10 mM sodium pyrophosphate, 10 mM NaF, 5 mM EDTA, 1 mM MgCl₂, 1 mM CaCl₂, 2 mM Na₃VO₄, 2 mM phenylmethylsulfonyl fluoride, 5 µg/mL aprotinin, 5 µg/mL leupeptin, 5 µg/mL pepstatin, 10% glycerol, and 1% Triton X-100). Homogenates were mixed by shaking for 45 min at 4°C and then centrifuged at 10,000xg for 20 min at 4°C. The supernatants were collected and the protein concentration was determined using BCA reagent (Pierce, Rockford, IL).

Samples were added to a reducing loading buffer with a final concentration of 50 mM Tris-HCl, pH 6.8, 10% glycerol, 2% SDS, 5% β-mercaptoethanol, and 0.005% bromophenol blue and heated for 10 min at 100°C. Equal amounts of protein were separated by SDS-PAGE (10% resolving gel), transferred to a nitrocellulose membrane and probed with either an anti-adiponectin rabbit polyclonal antibody (1:750) (Alpha Diagnostics, San Antonio, TX), anti-Ser₄₇₃ PKB rabbit polyclonal antibody (1:2000) (Cell Signaling Technology, Danvers, MA), anti-PKB (1:1000) (Cell Signaling Technology), or anti-AMPKα (1:1000) (Cell Signaling Technology) overnight at 4°C. Both the total and phospho-specific PKB antibodies detect all isoforms. The porcine adiponectin antibody was generated against the following two peptides: DQYQDKNVDQASGS and TEKPGALLPVPKGAC. Anti-adiponectin was purified from immunized rabbit serum by affinity chromatography. Membranes were then probed with a goat anti-rabbit IgG horseradish peroxidase (1:10,000) (Pierce) for 1 hour at room temperature. Blots were developed using the SuperSignal West Pico Chemiluminescent Substrate system (Pierce) and imaged with FOTO/Analyst Luminary FX system (Fotodyne Incorporated, Hartland, WI). Densitometry was performed using TotalLab TL100 software (TotalLab Ltd, Newcastle, UK). Serum (2 µL) was diluted in

reducing loading buffer, heat denatured, and separated by SDS-PAGE as described above for estimation of serum adiponectin concentrations. Phosphorylation of AMPK α at Thr₁₇₂ was measured by ELISA (Cell Signaling Technology).

Statistical Analysis. All statistical procedures outlined were performed using SAS 9.2 (SAS, Cary, NC). Residuals were analyzed to detect outliers and assess normality using the PROC Univariate procedure. Right-skewed data were log-transformed for further analysis. ANOVA was performed using the PROC GLM procedure. Main effects included diet and tissue where appropriate. Fixed effects include sex, group, and termination date. Fixed effects were used in the model when $p \leq 0.15$ for the fixed effect or interaction with main effect(s). Blood glucose data from the insulin challenge were analyzed using an autoregression order of one repeated measures. For flow cytometry data analysis was weighted on live cell number. Additionally, %CD203a⁺ was used as a covariate for analysis of CD14 and CD16 data. Data presented are least-squares means \pm standard error.

Results

Dietary fatty acid profiles are presented in **Table 3**. Only the LFC and HFPn3 diet contained measurable EPA and DHA while all three diets contained α -linolenic acid. The addition of n-3 PUFA to the HFPn3 diet decreased the n-6:n-3 from 47.6 in the HFP diet to 3.9 in the HFPn3 diet. The average calculated n-3 PUFA consumption was 0.40, 0, and 7.6 g/d for EPA, and 0.36, 0, and 6.7 g/d for DHA for the LFC, HFP, and HFPn3 groups, respectively. To maintain lean mature body weight, caloric intake was limited in the LFC group to an average of 1778 kcal/d compared to ~9800 kcal/day for the two high fat diets. Final body weights for the two high fat groups did not differ (**Table 4**).

To assess the impact of the high fat diet and n-3 supplementation on systemic metabolic disturbances, insulin, blood glucose, cholesterol, triglycerides, homocysteine, and blood pressure were measured (**Table 4**). There was no difference detected among the LFC, HFP, and HFPn3 groups in fasting serum insulin, cholesterol, triglyceride, or homocysteine concentrations. In contrast, there was a dietary effect with serum NEFA; the HFPn3 group had higher NEFA concentrations than the LFC group, whereas the HFP group was not different from either the LFC or the HFPn3 group. Additionally, consumption of either high fat diet increased systolic blood pressure, diastolic blood pressure, and mean arterial pressure. Although there were no differences in serum insulin concentrations, the LFC group had lower fasting blood glucose than either the HFP or HFPn3 groups. The high fasting blood glucose concentrations of pigs fed the HFP or HFPn3 diets caused an increase in HOMA-IR as compared with the LFC group.

Also summarized in **Table 4**, there were no differences in serum CRP or TNF α attributable to dietary treatment. However, serum CRP was positively correlated with fasting blood glucose ($r=0.42716$, $p=0.0421$). IL-6, IL-10, and IL-12 were detectable in the serum of only two pigs (data not shown). Serum adiponectin was decreased by both high fat diets vs. the control group, but the magnitude of this decrease was attenuated in pigs fed the HFPn3 diet (**Figure 1A**). Serum adiponectin was negatively correlated with fasting blood glucose ($r=-0.44268$, $p=0.0287$).

To assess whether differences in serum adiponectin were associated with altered AMPK activity, we measured LD muscle AMPK α phosphorylation. Both the HFP and HFPn3 groups had decreased phosphorylation of AMPK α at Thr₁₇₂, and a decreased ratio of Thr₁₇₂ AMPK α to total AMPK α as compared with the LFC group, but did not differ in total

AMPK α (**Figure 1B-D**). Phosphorylation of AMPK α at Thr₁₇₂ was positively correlated with serum adiponectin concentration ($r=0.47112$, $p=0.0233$) and negatively correlated with fasting blood glucose ($r=-0.49803$, $p=0.0156$). Additionally, we determined whether there was a dietary influence on ectopic lipid deposition. Percent lipid in liver and LD muscle was not influenced by diet (**Table 3**).

Insulin sensitivity was assessed by measuring the decrease in blood glucose following an insulin challenge. Blood glucose was decreased by 5 min in the LFC group, and by 10 min in the group fed the HFPn3 diet (**Figure 2**). The HFP group was unresponsive to insulin over the 15 min insulin challenge (**Figure 2**). Tissue differences in insulin signaling were assessed by evaluating phosphorylation of PKB at Ser₄₇₃. The HFPn3 diet decreased PKB abundance in SQ, VIS, and OM adipose depots (**Figure 3A, 3D, 3G**). However, there was no difference in Ser₄₇₃ phosphorylation of PKB (**Figure 3B, 3E, 3H**). The ratio of Ser₄₇₃ PKB to total PKB was decreased in the SQ and OM depots by the HFP diet, but not the HFPn3 diet (**Figures 3C, 3I**), but was unchanged in the VIS depot (**Figure 3F**). There was no difference in liver or LD muscle total PKB, or PKB phosphorylation (data not shown). Adipose tissue macrophages were evaluated by flow cytometry to determine both the extent of macrophage infiltration (CD203a) and phenotype. Phenotype was determined by the presence of CD16, an Fc γ receptor, and CD14, the co-receptor for tlr4. The percent of macrophages in adipose depot SVC was decreased in the HFP and HFPn3 groups with no difference amongst adipose depots (**Figure 4A**).

The HFP and HFPn3 groups had increased CD16⁺ macrophages in the adipose tissue compared with the LFC group, with no difference among the adipose depots (**Figure 4B**). There was no difference in CD16⁺ CD14⁺ macrophages among dietary treatments; however,

the OM adipose depot had decreased CD16⁺ CD14⁺ macrophages compared to the SQ adipose depot (**Figure 4C**). Additionally, the HFP diet increased the percent CD16⁺ CD14⁺ macrophages compared with the LFC and HFPn3 groups (**Figure 4D**) with no differences across adipose depots.

Discussion

Comparative animal models are critical to understanding the development of human obesity and the chronic inflammation in adipose tissue which contributes to the pre-diabetic state (i.e., metabolic syndrome). Although previous studies (11; 24; 33-34) have underscored the potential value of the Ossabaw pig as a model for obesity and pre-diabetes, this is to our knowledge the first study to address obesity-linked inflammation in adipose tissue of the Ossabaw pig, or the potential alleviation of this inflammation by dietary n-3 fatty acids. Whereas others (20) have reported increased circulating concentrations of TNF α in response to diet-induced obesity in Ossabaw swine, we found no diet-related increase in serum concentrations of common pro-inflammatory markers, TNF α , CRP, IL-6, or IL-12, despite the prolonged duration of our study and the marked development of obesity (i.e., over 100 kg weight gained beyond mature body weight). Whether this difference relates to differences in diet composition across studies is currently unresolved. It is noteworthy that whereas others (20) have attributed both inflammation and liver injury to the presence of fructose in atherogenic diets, our diet contained sucrose which contributed 14% fructose to the diet (on a weight basis), and no added cholesterol. However, it is also intriguing to note that reduced ectopic storage of lipid in liver and skeletal muscle due to increased capacity for expansion of adipose mass has been associated with improvements in metabolic markers associated with obesity (17), and presumably, alleviation of obesity-linked inflammation. We saw no

evidence of ectopic lipid accumulation in the liver or skeletal muscle of pigs fed either high fat diet. Consequently, the marked capacity of the Ossabaw swine used in our study to expand adipose mass may in fact afford them some protection against obesity-linked inflammation.

To evaluate adipose inflammation in this model and make comparisons with human literature, we investigated the adipose tissue macrophage population in lean vs. obese Ossabaw swine. Several studies of human adipose tissue have categorized macrophages as being CD14⁺ (4; 18; 35; 41). In our study, only a small percentage (8-10%) of the adipose-derived SVC were identified as CD14⁺. Consequently, we used the mature macrophage marker, CD203a, to identify macrophage populations and the markers CD14 and CD16 to assess phenotypic changes associated with diet or adipose depot. Surprisingly, both the HFP and HFPn3 groups had fewer macrophages than the LFC group, although this decrease did not occur in all depots across diets. It seems possible that accelerated macrophage turnover in obese pigs led to an increase in adipose monocytes not expressing CD203a. It is also possible that recruitment of adipogenic progenitors, due to rapid expansion of adipose mass in pigs fed the high fat diets, decreased the overall percentage of macrophages in the SVC population.

The predominant population of adipose macrophages in the Ossabaw pig were CD16⁺, with the vast majority of these being CD14⁻. This is quite intriguing as there is a paucity of information relating to this particular phenotype of adipose macrophages. Given that CD16⁺CD14⁻ monocytes are anti-inflammatory (7), it is quite possible that adipose inflammation in obese Ossabaw swine may be minimized by the majority of macrophages being of the CD16⁺CD14⁻ phenotype. However, it is important to note that there was in fact

an increase in the number of CD16⁻CD14⁺ macrophages detected in the adipose tissue of pigs fed the HFP diet vs. the LFC or HFPn3 groups. In humans, classical CD16⁻CD14⁺ blood monocytes express high levels of CCR2 and CD62L (29), and are thus pro-inflammatory (7). This perhaps indicates that the greater number of CD16⁻CD14⁺ macrophages in the adipose tissue of pigs fed the HFP diet reflects obesity-linked infiltration with pro-inflammatory cells, and that the HFPn3 diet protected against this infiltration, despite the marked obesity. It is noteworthy, that we have previously demonstrated that dietary supplementation with n-3 fatty acids causes enrichment of both adipose tissue and muscle (14), and thus higher concentrations of n-3 PUFA in the adipose tissue of pigs fed the HFPn3 diet may have protected them from the increase in the CD16⁻CD14⁺ macrophages.

As regards indicators of pre-diabetes, fasting hyperglycemia was readily induced by both high fat diets, and this occurred with increased HOMA-IR and blood pressure measures. Hyperglycemia and hyperinsulinemia have been reported (19-20) in pig models, although insulin peaked at one month and decreased thereafter in one study (19). The degree of hyperglycemia induced by the high fat diets without concomitant hyperinsulinemia is striking. There are instances where hypoinsulinemia is present in a diabetic state (9), and may be caused decreased beta cell function due to chronic hyperglycemia (10). However, this has not been addressed in this model to our knowledge. Although, cholesterol and triglycerides have also been increased by high fat diets in several studies utilizing swine (11; 19-20; 24), the diets used in our study did not cause significant increases in either. It should be noted that we did not add cholesterol to the diets as other researchers have done (11; 20; 24). In Ossabaw swine, dietary cholesterol likely plays a pivotal role in the development of hypercholesterolemia and hypertriglyceridemia. Circulating NEFA were increased in only in

the HFPn3 group vs. the LFC group. This is most likely a consequence of the slightly higher fat content of the HFPn3 diet due to addition of the n3 fatty acid source.

In the context of obesity, the activity of AMP-activated protein kinase (AMPK) is quite important due to its stimulation of glucose uptake and fatty acid oxidation in muscle (31; 40). Relative to the control group, we found that both obese groups had decreased abundance of phosphorylated Thr₁₇₂ AMPK α and a decrease in the ratio of phosphoThr₁₇₂ AMPK α to total AMPK α in the LD muscle. Genetically, Ossabaw swine contain the mutation Val₁₉₉→Ile in the PRKAG3 (the γ_3 isoform of AMPK) gene which negatively affects the activity of AMPK (22). It has been hypothesized that decreased AMPK activity due to this mutation is causal in the development of insulin resistance in Ossabaw pigs (27). Although adiponectin clearly stimulates activation of AMPK (40), it seems unlikely that the relatively small decrease in serum adiponectin in the obese groups vs. the control group contributed to the reduction in Thr₁₇₂ AMPK α . Given the PRKAG3 mutation, it will indeed be intriguing to determine whether Ossabaw pigs are more susceptible to the effects of high fat diets on AMPK activity than are pigs of commercial lean lines.

Changes in blood glucose concentrations during the insulin challenge indicated that n-3 fatty acid supplementation improved insulin sensitivity vs. pigs fed the HFP diet. However, there was no dietary difference in Ser₄₇₃ PKB in adipose tissue, liver, or LD muscle after insulin stimulation. Total PKB was decreased in all adipose depots compared to the LFC group and Ser₄₇₃ PKB:PKB was increased in SQ and OM adipose depots in pigs fed the HFPn3 diet compared to the HFP pigs. In adipocytes, activation of peroxisome proliferator-activated receptor (PPAR) γ by thiazolidinediones decreases PKB α abundance specifically (38). Previous work has shown that n3 fatty acids (or their metabolites) are

ligand activators of PPAR γ (3; 25) and it is quite possible that the reduction in PKB reflects this activity of n-3 fatty acids in adipocytes. Nonetheless, the reduction in PKB did not preclude the stimulation of glucose uptake by insulin during the insulin challenge. Additionally, it should be noted that the insulin challenge was performed under anesthesia, which although necessary for this study, is also limitation.

In summary, we have shown marked obesity, hyperglycemia, hypertension and a modest reduction in circulating adiponectin concentrations in Ossabaw swine fed a diet high in saturated fat. However, there were only limited indications of the expected obesity-linked inflammation in adipose tissue. The predominate evidence of inflammation was the increased CD16⁺CD14⁺ macrophages in the adipose tissue of the HFP group, which are most likely pro-inflammatory macrophages. Despite the lack of a clear inflammatory response, the HFP group had decreased sensitivity to administration of exogenous insulin as compared with the control and HFPn3 groups. Major effects of n-3 fatty acid supplementation were determined to be the regulation of PKB abundance, improved insulin-stimulated glucose clearance during the insulin challenge, and blunted infiltration of adipose tissue with CD16⁺CD14⁺ macrophages. Further investigation with the monocytic markers CD172a and CD163 may provide further insight to adipose depot macrophage populations in this novel model of obesity and pre-diabetes. Finally, the loss of three pigs from the HFPn3 dietary treatment, although not definitively linked to diet, limited our statistical power in studying the beneficial effects of n-3 PUFA in this model.

Acknowledgements

This project was supported by USDA National Research Initiative Grant 2008-01847 from the USDA National Institute for Food and Agriculture.

References

1. Ajuwon KM, Spurlock ME. 2005. Palmitate activates the NF-kappaB transcription factor and induces IL-6 and TNFalpha expression in 3T3-L1 adipocytes. *J Nutr* 135:1841-1846.
2. Bahceci M, Gokalp D, Bahceci S, Tuzcu A, Atmaca S, Arikan S. 2007. The correlation between adiposity and adiponectin, tumor necrosis factor alpha, interleukin-6 and high sensitivity C-reactive protein levels. Is adipocyte size associated with inflammation in adults? *J Endocrinol Invest* 30:210-214.
3. Banga A, Unal R, Tripathi P, Pokrovskaya I, Owens RJ, Kern PA, Ranganathan G. 2009. Adiponectin translation is increased by the PPARgamma agonists pioglitazone and omega-3 fatty acids. *Am J Physiol Endocrinol Metab* 296:E480-489.
4. Bourlier V, Zakaroff-Girard A, Miranville A, De Barros S, Maumus M, Sengenès C, Galitzky J, Lafontan M, Karpe F, Frayn KN, Bouloumie A. 2008. Remodeling phenotype of human subcutaneous adipose tissue macrophages. *Circulation* 117:806-815.
5. Bray GA, Lovejoy JC, Smith SR, DeLany JP, Lefevre M, Hwang D, Ryan DH, York DA. 2002. The influence of different fats and fatty acids on obesity, insulin resistance and inflammation. *J Nutr* 132:2488-2491.
6. Cordain L, Eaton SB, Sebastian A, Mann N, Lindeberg S, Watkins BA, O'Keefe JH, Brand-Miller J. 2005. Origins and evolution of the Western diet: health implications for the 21st century. *Am J Clin Nutr* 81:341-354.
7. Cros J, Cagnard N, Woollard K, Patey N, Zhang SY, Senechal B, Puel A, Biswas SK, Moshous D, Picard C, Jais JP, D'Cruz D, Casanova JL, Trouillet C, Geissmann F. 2010. Human CD14^{dim} monocytes patrol and sense nucleic acids and viruses via TLR7 and TLR8 receptors. *Immunity* 33:375-386.
8. Davis JE, Gabler NK, Walker-Daniels J, Spurlock ME. 2008. Tlr-4 deficiency selectively protects against obesity induced by diets high in saturated fat. *Obesity (Silver Spring)* 16:1248-1255.
9. DeFronzo RA. 2004. Pathogenesis of type 2 diabetes mellitus. *Med Clin North Am* 88:787-835, ix.
10. Donath MY, Halban PA. 2004. Decreased beta-cell mass in diabetes: significance, mechanisms and therapeutic implications. *Diabetologia* 47:581-589.

11. Dyson MC, Alloosh M, Vuchetich JP, Mokelke EA, Sturek M. 2006. Components of metabolic syndrome and coronary artery disease in female Ossabaw swine fed excess atherogenic diet. *Comp Med* 56:35-45.
12. Flachs P, Mohamed-Ali V, Horakova O, Rossmeisl M, Hosseinzadeh-Attar MJ, Hensler M, Ruzickova J, Kopecky J. 2006. Polyunsaturated fatty acids of marine origin induce adiponectin in mice fed a high-fat diet. *Diabetologia* 49:394-397.
13. Folch J, Lees M, Sloane Stanley GH. 1957. A simple method for the isolation and purification of total lipides from animal tissues. *J Biol Chem* 226:497-509.
14. Gabler NK, Spencer JD, Webel DM, Spurlock ME. 2008. n-3 PUFA attenuate lipopolysaccharide-induced down-regulation of toll-like receptor 4 expression in porcine adipose tissue but does not alter the expression of other immune modulators. *J Nutr Biochem* 19:8-15.
15. Itoh M, Suganami T, Satoh N, Tanimoto-Koyama K, Yuan X, Tanaka M, Kawano H, Yano T, Aoe S, Takeya M, Shimatsu A, Kuzuya H, Kamei Y, Ogawa Y. 2007. Increased adiponectin secretion by highly purified eicosapentaenoic acid in rodent models of obesity and human obese subjects. *Arterioscler Thromb Vasc Biol* 27:1918-1925.
16. Kalupahana NS, Claycombe K, Newman SJ, Stewart T, Siriwardhana N, Matthan N, Lichtenstein AH, Moustaid-Moussa N. 2010. Eicosapentaenoic acid prevents and reverses insulin resistance in high-fat diet-induced obese mice via modulation of adipose tissue inflammation. *J Nutr* 140:1915-1922.
17. Kim JY, van de Wall E, Laplante M, Azzara A, Trujillo ME, Hofmann SM, Schraw T, Durand JL, Li H, Li G, Jelicks LA, Mehler MF, Hui DY, Deshaies Y, Shulman GI, Schwartz GJ, Scherer PE. 2007. Obesity-associated improvements in metabolic profile through expansion of adipose tissue. *J Clin Invest* 117:2621-2637.
18. Kovacikova M, Sengenès C, Kovacova Z, Siklova-Vitkova M, Klimcakova E, Polak J, Rossmeislova L, Bajzova M, Hejnova J, Hnevkovska Z, Bouloumie A, Langin D, Stich V. 2010. Dietary intervention-induced weight loss decreases macrophage content in adipose tissue of obese women. *Int J Obes (Lond)*.
19. Lee J, Xu Y, Lu L, Bergman B, Leitner JW, Greyson C, Draznin B, Schwartz GG. 2010. Multiple abnormalities of myocardial insulin signaling in a porcine model of diet-induced obesity. *Am J Physiol Heart Circ Physiol* 298:H310-319.
20. Lee L, Alloosh M, Saxena R, Van Alstine W, Watkins BA, Klaunig JE, Sturek M, Chalasani N. 2009. Nutritional model of steatohepatitis and metabolic syndrome in the Ossabaw miniature swine. *Hepatology* 50:56-67.

21. Lepage G, Roy CC. 1986. Direct transesterification of all classes of lipids in a one-step reaction. *J Lipid Res* 27:114-120.
22. Lloyd PG, Fang M, Brisbon IL, Andersson L, Sturek M. 2006. AMP kinase gene mutation is consistent with a thrifty phenotype (metabolic syndrome) in a population of feral swine. *FASEB J* 20:A299.
23. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. 1985. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412-419.
24. Neeb ZP, Edwards JM, Alloosh M, Long X, Mokelke EA, Sturek M. 2010. Metabolic syndrome and coronary artery disease in Ossabaw compared with Yucatan swine. *Comp Med* 60:300-315.
25. Oster RT, Tishinsky JM, Yuan Z, Robinson LE. 2010. Docosahexaenoic acid increases cellular adiponectin mRNA and secreted adiponectin protein, as well as PPARgamma mRNA, in 3T3-L1 adipocytes. *Appl Physiol Nutr Metab* 35:783-789.
26. Satoh N, Shimatsu A, Kotani K, Sakane N, Yamada K, Suganami T, Kuzuya H, Ogawa Y. 2007. Purified eicosapentaenoic acid reduces small dense LDL, remnant lipoprotein particles, and C-reactive protein in metabolic syndrome. *Diabetes Care* 30:144-146.
27. Shen W, Sturek M, Liechty EA. 2007. Molecular mechanisms for insulin resistance in Ossabaw miniature swine. *FASEB J* 21:A1053.
28. Shi H, Kokoeva MV, Inouye K, Tzameli I, Yin H, Flier JS. 2006. TLR4 links innate immunity and fatty acid-induced insulin resistance. *J Clin Invest* 116:3015-3025.
29. Tacke F, Randolph GJ. 2006. Migratory fate and differentiation of blood monocyte subsets. *Immunobiology* 211:609-618.
30. Todoric J, Loffler M, Huber J, Bilban M, Reimers M, Kadl A, Zeyda M, Waldhausl W, Stulnig TM. 2006. Adipose tissue inflammation induced by high-fat diet in obese diabetic mice is prevented by n-3 polyunsaturated fatty acids. *Diabetologia* 49:2109-2119.
31. Tomas E, Tsao TS, Saha AK, Murrey HE, Zhang Cc C, Itani SI, Lodish HF, Ruderman NB. 2002. Enhanced muscle fat oxidation and glucose transport by ACRP30 globular domain: acetyl-CoA carboxylase inhibition and AMP-activated protein kinase activation. *Proc Natl Acad Sci U S A* 99:16309-16313.
32. Tsukumo DM, Carvalho-Filho MA, Carvalheira JB, Prada PO, Hirabara SM, Schenka AA, Araujo EP, Vassallo J, Curi R, Velloso LA, Saad MJ. 2007. Loss-of-function

- mutation in Toll-like receptor 4 prevents diet-induced obesity and insulin resistance. *Diabetes* 56:1986-1998.
33. Wangsness PJ, Martin RJ, Gahagan JH. 1977. Insulin and growth hormone in lean and obese pigs. *Am J Physiol* 233:E104-108.
 34. Wangsness PJ, Martin RJ, Gatchel BB. 1980. Insulin induced growth hormone response in fast-growing, lean and in slow-growing, obese pigs. *Growth* 44:318-326.
 35. Wentworth JM, Naselli G, Brown WA, Doyle L, Phipson B, Smyth GK, Wabitsch M, O'Brien PE, Harrison LC. 2010. Pro-inflammatory CD11c+CD206+ adipose tissue macrophages are associated with insulin resistance in human obesity. *Diabetes* 59:1648-1656.
 36. Williams KT, Garrow TA, Schalinske KL. 2008. Type I diabetes leads to tissue-specific DNA hypomethylation in male rats. *J Nutr* 138:2064-2069.
 37. Xi S, Yin W, Wang Z, Kusunoki M, Lian X, Koike T, Fan J, Zhang Q. 2004. A minipig model of high-fat/high-sucrose diet-induced diabetes and atherosclerosis. *Int J Exp Pathol* 85:223-231.
 38. Xiao Y, Yuan T, Yao W, Liao K. 2010. 3T3-L1 adipocyte apoptosis induced by thiazolidinediones is peroxisome proliferator-activated receptor- γ -dependent and mediated by the caspase-3-dependent apoptotic pathway. *FEBS J* 277:687-696.
 39. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS, Tartaglia LA, Chen H. 2003. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest* 112:1821-1830.
 40. Yamauchi T, Kamon J, Minokoshi Y, Ito Y, Waki H, Uchida S, Yamashita S, Noda M, Kita S, Ueki K, Eto K, Akanuma Y, Froguel P, Foufelle F, Ferre P, Carling D, Kimura S, Nagai R, Kahn BB, Kadowaki T. 2002. Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med* 8:1288-1295.
 41. Zeyda M, Farmer D, Todoric J, Aszmann O, Speiser M, Gyori G, Zlabinger GJ, Stulnig TM. 2007. Human adipose tissue macrophages are of an anti-inflammatory phenotype but capable of excessive pro-inflammatory mediator production. *Int J Obes (Lond)* 31:1420-1428.

Tables

Table 2. *Ingredient composition of the diet (%) as-fed.*

Ingredient	Diet		
	LFC	HFP	HFPn3
Corn	46.92	25.06	13.32
Soybean Meal-48	4.00	19.55	21.35
Sucrose	-	28.00	28.00
Palm Oil	-	18.00	18.00
Menhaden Meal	7.50	-	-
Soybean Hulls	33.44	-	-
Wheat Bran	3.00	-	-
Canola Oil	1.00	-	3.00
Gromega TM 365 ¹	1.00	-	5.00
Meat & Bone Meal ²	-	6.25	6.25
Vitamin Mix	0.50	0.50	0.50
Salt	0.40	0.40	0.40
Mineral Mix	0.10	0.10	0.10
Selenium Mix	0.05	0.05	0.05
DL-Methionine	-	0.07	0.09
Dicalcium Phosphate	1.88	1.63	1.64
Limestone	0.21	0.38	2.30
Ethoxyquin ³	0.0125	0.0125	0.0125
Metabolizable Energy (kcal/kg)	2750	4059	4058
Fat (kcal/kg)	382	1572	1877
Ether Extract (%)	4.69	20.27	23.98
Crude Protein (%)	14.92	15.14	15.00
Lysine (%)	0.81	0.80	0.83
Methionine (%)	0.29	0.29	0.29
Met + Cys (%)	0.47	0.50	0.50
Crude Fiber (%)	13.69	1.39	1.19
Calcium (%)	0.90	1.20	1.93
Available Phosphorous	0.60	0.60	0.60

¹JBS United, Sheridan, IN; ²Darling International, Des Moines, IA; ³Novus International, St.

Table 3. *Dietary fatty acid profile (g/100 g total fatty acid).*

Fatty Acid	Diet		
	LFC	HFP	HFPn3
6:0	0.00	0.00	0.00
8:0	0.00	0.00	0.00
10:0	0.66	0.80	0.75
12:0	0.00	1.44	2.12
14:0	2.34	1.06	1.59
14:1	0.00	0.00	0.00
16:0	18.47	43.98	38.85
16:1	2.51	0.00	0.89
18:0	4.08	5.18	4.83
18:1	28.14	36.6	37.58
18:2n6	34.14	10.73	10.49
18:3n6	0.00	0.00	0.00
18:3n3	3.77	0.23	1.03
20:0	0.44	0.00	0.21
20:1	0.66	0.00	0.00
20:2	0.00	0.00	0.00
20:3	0.00	0.00	0.00
20:3n6	0.00	0.00	0.00
20:4n6	0.00	0.00	0.00
20:5n3	2.33	0.00	0.88
22:1	0.00	0.00	0.00
22:2	0.00	0.00	0.00
24:0	0.00	0.00	0.00
22:6n3	2.10	0.00	0.78
Other	0.37	0.00	0.00
Total	100.00	100.00	100.00
Saturated	26.36	52.45	48.35
n3	8.20	0.23	2.69
n6	34.14	10.73	10.49
n6/n3	4.16	47.59	3.91

Table 4. *Growth and metabolic parameters of Ossabaw swine at termination.*

Parameter	Diet			p-value
	LFC	HFP	HFPn3	
Final Body Weight (kg)	49.6±4.7 ^a	156.7±5.3 ^b	154.4±5.8 ^b	<0.0001
Insulin (pg/mL)	63.5±8.6	80.3±11.7	76.5±14.7	0.44
Blood Glucose (mg/dL)	119.8±14.8 ^a	233.5±29.4 ^b	221.5±34.4 ^b	0.0011
HOMA-IR	0.44±0.12 ^a	1.03±0.13 ^b	0.97±0.18 ^b	0.0121
Cholesterol (mg/dL)	68.4±5.7	90.8±6.4	79.0±6.9	0.06
Triglycerides (mg/dL)	42.0±7.1	61.9±7.4	61.4±8.7	0.11
Liver Lipid (%)	4.23±0.18	4.54±0.18	4.58±0.22	0.39
LD Lipid (%)	3.53±1.06	5.90±1.10	5.98±1.29	0.21
NEFA (mM)	0.57±0.22 ^a	1.13±0.23 ^{ab}	1.46±0.27 ^b	0.0453
CRP (ng/mL)	13.7±2.5	20.6±2.6	18.4±3.0	0.17
TNFα (pg/mL)	24.9±3.2	21.6±2.8	15.5±2.3	0.10
Homocysteine (μM)	11.5±5.8	33.0±5.2	26.2±8.1	0.06
Diastolic BP	63.4±7.9 ^a	96.3±7.1 ^b	102.0±9.0 ^b	0.0251
Systolic BP	127.2±6.9 ^a	170.1±6.3 ^b	178.7±8.1 ^b	0.0021
Mean Arterial Pressure	93.6±8.9 ^a	118.0±7.9 ^b	134.0±10.2 ^b	0.0275

Figures




Figure 1. *n-3 fatty acids attenuate the high palm oil induced decrease in serum adiponectin, but do not alter AMPK.*

(A) Relative serum adiponectin and (B) AMPK α , (C) Thr₁₇₂ AMPK α , and (D) Thr₁₇₂ AMPK α :AMPK α of LD Ossabaw swine fed low fat control (LFC), high fat palm oil (HFP), or high fat palm oil plus n-3 fatty acids (HFPn3) diets. Values are (A-B) arbitrary densitometric units or (C) arbitrary units per mg protein.. Means with different letters represent a difference of $p < 0.05$. Main effects (A) Diet $p = 0.0410$; (B) Diet $p = 0.42$; (C) Diet $p = 0.0055$; (D) Diet $p = 0.0136$.




Figure 2. *n-3 fatty acids attenuate high palm oil induced insulin resistance.*

Blood glucose from Ossabaw swine challenged with 0.25 IU/kg porcine insulin after a 12 hour fast from 3 dietary treatments: low fat control (LFC), high fat palm oil (HFP), high fat palm oil plus n-3 fatty acids (HFPn3). Differences are from 0 to 5 min, 5 to 10 min, and 10 to 15 min within diet are represented by *, †, and ‡, respectively. Main effects: Diet $p=0.0004$, Time $p<0.0001$, Diet*Time $p=0.0481$.




Figure 3. *Insulin resistance is not reflected in adipose tissue S₄₇₃ PKB phosphorylation.*

(A) PKB, (B) Ser₄₇₃ PKB, and (C) Ser₄₇₃ PKB:PKB of subcutaneous adipose (SQ), (D) PKB, (E) Ser₄₇₃ PKB, and (F) Ser₄₇₃ PKB:PKB of visceral adipose (VIS), and (G) PKB, (H) Ser₄₇₃ PKB, and (I) Ser₄₇₃ PKB:PKB of omental adipose (OM) from Ossabaw swine fed low fat control (LFC), high fat palm oil (HFP), or high fat palm oil plus n-3 fatty acids (HFPn3) diets. Values are arbitrary densitometric units. Means with different letters represent difference of $p < 0.05$. Main effects: (A) Diet $p = 0.0238$; (B) Diet $= 0.34$; (C) Diet $p = 0.0141$; (D) Diet $p = 0.0326$; (E) Diet $= 0.18$; (F) Diet $p = 0.22$; (G) Diet $p = 0.0153$; (H) Diet $= 0.12$; (I) Diet $p = 0.0032$.




Figure 4. High fat diets have fewer macrophages, but phenotype is altered by obesity and depot.

(A) Macrophages (CD203a), and (B) CD16⁺ macrophages by diet, (C) CD16⁺CD14⁺ macrophages by adipose depot, and (D) CD16⁻ CD14⁺ macrophages by diet in subcutaneous (SQ), visceral (VIS), and omental (OM) adipose depots from Ossabaw swine fed low fat control (LFC), high fat palm oil (HFP), or high fat palm oil plus n-3 fatty acids (HFPn3) diets. Values are percent positive out of SVC's. Means with different letters represent difference of $p < 0.05$. Main effects: (A) Diet $p = 0.0002$, Tissue $p = 0.9192$, Diet*Tissue $p = 0.0007$; (B) Diet $p = p < 0.0001$, Tissue $p = 0.10$, Diet*Tissue $p = 0.21$; (C) Diet $p = 0.50$, Tissue $p < 0.0443$, Diet*Tissue $p = 0.98$; (D) Diet $p = 0.0053$, Tissue $p = 0.65$, Diet*Tissue $p = 0.74$.

CHAPTER 4: ADIPOGENIC POTENTIAL OF OSSABAW ADIPOCYTES PROTECTS AGAINST IL-8 SECRETION

Abstract

Some obese individuals appear to be protected against obesity-linked adipose tissue inflammation and the ensuing metabolic consequences. This protection may be afforded by an inherent ability to markedly expand adipose tissue by hyperplasia in response to excess caloric intake, as it has been previously demonstrated that adipose tissue expansion improves both the metabolic and inflammatory profile. It was determined that Ossabaw swine develop minimal evidence of inflammation, despite morbid obesity induced by long-term consumption of a high fat diet. The ability of Ossabaw swine to store large amounts of lipid in adipose tissue when provided excess calories may have provided protection against the development of inflammation. Thus, we hypothesized that Ossabaw adipocytes will have increased adipogenesis and be protected against palmitate-induced inflammation and insulin resistance compared with Yorkshire adipocytes. Ossabaw adipocytes have increased clonal expansion, PPAR γ expression, and G3PDH activity compared to Yorkshire adipocytes. Palmitate activated NF κ B, but activation was not different between Ossabaw and Yorkshire adipocytes. However, palmitate only induced IL-8 production in Yorkshire adipocytes. The increased adipogenic potential of Ossabaw adipocytes may be protective against palmitate induced IL-8 production. This is confirmed by the lack of obesity-linked increases in serum and adipose tissue IL-8 concentrations in Ossabaw swine. As determined by 2-NBDG uptake, insulin sensitivity was decreased by both palmitate, irrespective of genotype. However, Ossabaw adipocytes had both lower basal and insulin-stimulated 2-NBDG uptake.

Thus, the *in vitro* evidence supports that the adipogenic potential of Ossabaw adipocytes may be protective against palmitate induced inflammation with respect to IL-8 production.

Introduction

The prevalence of obesity is increasing in the United States (16). Obesity is considered a state of chronic low-grade inflammation. Nearly all of the co-morbidities associated with obesity can be linked to inflammation (3; 29). It is estimated that ~80% of individuals with type 2 diabetes (31) and 65% of individuals with the metabolic syndrome (15) are obese. Some obese individuals appear to be protected against obesity-linked inflammation and metabolic disturbances (22; 38). These individuals are termed metabolically healthy but obese (MHO). MHO individuals have decreased inflammation compared to their un-healthy counterparts, including lower C-reactive protein (CRP) (21; 37), interleukin 6 (IL-6) (24; 37), and tumor necrosis factor (TNF) α (24) as well as increased adiponectin (1; 44).

In addition to decreased inflammation, MHO individual have a more favorable metabolic profile. Compared with their un-healthy obese counterparts, MHO individuals have less visceral fat (7; 18; 20-21; 24; 44) and less ectopic lipid deposition in muscle (39; 44) and liver (39), despite similar BMI (7; 18; 20-21; 37) and body fat percentage (7; 21). Additionally, MHO individuals have decreased circulating concentrations of triglycerides (7; 18; 20-21; 24; 37), total cholesterol (24), LDL-cholesterol (24; 37), and increased concentrations of HDL-cholesterol (7; 21; 24; 37) compared to their un-healthy counterparts. MHO individuals retain insulin sensitivity (7; 18; 20-21; 37; 39) and have lower fasting insulin (7; 20-21) and glucose (7; 18; 20).

An extraordinary ability to expand subcutaneous adipose tissue may be a fundamental component of the favorable metabolic profile exhibited by MHO individuals. Evidence suggests that adipose tissue has a genetic limit on its ability to expand by hypertrophy and hyperplasia; and that the ability to expand varies by depot (12). Excess energy intake may exhaust the body's ability to store fat in adipose tissue, leading to ectopic lipid deposition in the muscle and liver. Ectopic lipid deposition has been closely linked to the development of insulin resistance (45). In *ob/ob* mice (devoid of leptin), the induction of adipose tissue expansion by over-expression of adiponectin attenuates insulin resistance and improves systemic inflammation normally seen in this model, despite morbid obesity (23).

Swine are increasingly being used as a biomedical model for researching obesity, insulin resistance, and cardiovascular disease (8; 10; 26-27; 30; 35). Ossabaw swine are an interesting model for studying obesity and its co-morbidities due to the breed's ability to accumulate lipid, develop impaired glucose tolerance, and frequent development of insulin resistance (28; 30; 42-43). However, we recently found that when fed a high fat diet for 30 weeks, Ossabaw swine display little evidence of the expected chronic low-grade inflammation associated with morbid obesity (14). Potentially, the ability of Ossabaw swine to accumulate adipose tissue may protect them against the development of inflammation. Thus, we hypothesized that the ability of Ossabaw swine to expand adipose tissue more readily than lean commercial swine will be evident by increased adipogenesis in Ossabaw stromal vascular cell (SVC) derived adipocytes and that they will have decreased palmitate-induced inflammation and insulin resistance compared to Yorkshire SVC derived adipocytes.

Methods

Animals. All animal procedures were conducted at Iowa State University and were approved by the Iowa State University Institutional Animal Care and Use Committee. To demonstrate the ability of Ossabaw swine to expand adipose tissue, subcutaneous (SQ) and visceral (VIS) adipose tissue was collected from a study previously described (14). Briefly, 27 mature male and female Ossabaw swine were fed three diets for 30 weeks: a restricted Low Fat Control (LFC), an *ad libitum* High Fat Palm Oil (HFP), and an *ad libitum* High Fat Palm Oil plus n-3 fatty acids (HFPn3). Final body weights for the three diets were 49.6, 156.7 and 154.4 kg, respectively. Additionally, SQ adipose tissue was collected from Yorkshire swine (~110 kg) to compare adipocyte size.

Adipocyte Size. Frozen sections of adipose depots were placed in 10% neutral buffered formalin overnight and then embedded in paraffin. Thereafter, sections of 5 μ m thickness were mounted onto microscope slides for staining with hematoxylin and eosin for analysis of adipocyte area and adipocyte number. Slides were imaged with a LEICA DM1 3000B microscope (LEICA Microsystems Inc., Buffalo Grove, IL) equipped with a QICAM Fast 1394 camera (QIMAGING, Surrey, BC, Canada). Adipocyte area was analyzed from an average of 6 fields at 10X magnification with AxioVision LE 4.8.2 (Carl Zeiss AG, Thornwood, NY). Measurements were taken from duplicate slides.

Cell Culture. SVC were isolated from lean Ossabaw and Yorkshire swine as previously described (14). Cells were grown in LG-DMEM (pH 7.4) supplemented with 10% FBS, 100 U/mL penicillin, 100 μ g/mL Streptomycin, 5 μ g/mL gentamicin, and 2.5 μ g/mL Amphotericin B. Media was changed every two days during the growth phase until one day after confluency. For differentiation of SVC into adipocytes, growth media was

supplemented with 100 nM insulin, 0.1% ITS (Sigma-Aldrich, St. Louis, MO), 0.5 mM 3-isobutyl-1-methylxanthine, 1 μ M dexamethasone, 33 μ M biotin, 17 μ M pantothenic acid, and 5 μ M troglitazone for 6 days. Thereafter, the media was changed at days 6 and 8 to growth media supplemented with 100 nM insulin 0.25 μ M dexamethasone, 33 μ M biotin, and 17 μ M pantothenic acid. All experiments utilized SVC between passages 3 and 6.

SVC Proliferation and Differentiation. SVC growth phase proliferation was determined for 3 days using BrDU incorporation (Millipore, Billerica, MA). Additionally, clonal expansion was measured for the first 3 days of differentiation using BrDU incorporation (Millipore). At days 10 days post-differentiation, adipocytes were stained with Oil Red O as follows: Cells were fixed for one hour with 10% neutral buffered formalin. Then were washed 3X with 60% isopropanol and stained for 10 minutes in a 0.21% w/v Oil Red-O in 60% isopropanol. Cells were then washed 4X with tap water and covered with 50% glycerol. Images were taken as described above for quantification of lipid droplet size. Cells were then washed 4X with tap water and air dried. Oil Red O was extracted with 100% isopropanol for 30 minutes with rocking and was quantified by measuring the absorbance at 500 nm.

G3PDH. The enzyme activity of glycerol-3-phosphate dehydrogenase (G3PDH) was determined from 10 day post-differentiation Ossabaw and Yorkshire SVC cell lysate. The assay for G3PDH was performed as described by Gamou et al. (17). Briefly, cell lysate was added to 100 μ L of reaction buffer (125 mM Triethanolamine-HCl pH 7.5, 2.5 mM EDTA, 0.5 mM NADH, 1.1 mM dihydroxyacetone phosphate, 0.125 mM β -mercaptoethanol) and the absorbance was read at 340 nm. Reagents for G3PDH activity were obtained from

Sigma-Aldrich (St. Louis, MO). Absorbance measurements were taken at 340 nm every 30 s for 7 min at 25°C on Synergy 4 microplate reader (Bio-Tek, Winooski, VT). G3PDH activity was calculated from the change in absorbance at 340 nm and expressed as nmol NAD(P)H per min per mg protein.

Insulin Sensitivity. Ossabaw and Yorkshire adipocytes (10 day post differentiation) were treated with Vehicle (untreated), 0.5 mM sodium palmitate, or 10 µg/mL LPS (*Escherichia coli* serotype O55:B55) in LG-DMEM (pH 7.4) with 0.1 mM BSA, 100 U/mL penicillin, 100 µg/mL Streptomycin for 48 hours with a media change at 24 hours. At 48 hours, cells were washed once with PBS, and then incubated with or without 100 nM insulin for 15 minutes. Thereafter 2-NBDG (2-deoxy-[(7-nitro-2,1,3-benzoxadiazol-4-yl)amino]-D-glucose) (Cayman Chemical, Ann Arbor, MI) was added at a final concentration of 100 µg/mL and incubated for 30 minutes. Cells were washed twice with PBS and the uptake of 2-NBDG was measured by fluorescence with the excitation/emission at 485/535 nm. Cellular protein was collected and protein was measured using the BCA reagent (Pierce, Rockford, IL). Total cellular protein was used as a covariate for 2-NBDG uptake.

Western Blot Analysis. Western blots were performed as previously described (14) with protein isolated from palmitate and LPS treated adipocytes. Nitrocellulose membranes were probed with PPAR γ rabbit polyclonal antibody (Millipore).

ELISA. Nuclear Factor κ B p 65 nuclear translocation was measured on palmitate and LPS treated cells using an ELISA (Cayman Chemical). Nuclear Extracts were extracted according to Cayman Chemical protocol with modifications. Protease and phosphatase inhibitors used in the extractions were 2 mM Na₃VO₄, 2 mM phenylmethylsulfonyl fluoride,

5 µg/mL aprotinin, 5 µg/mL leupeptin, 5 µg/mL pepstatin for both the hypotonic buffer and the nuclear extraction buffer. Media IL-8 and IL-6 were measured on 48 hour cell culture treatment media (R&D Systems, Minneapolis, MN). Additionally, IL-8 concentration was determined on serum and SQ and VIS adipose tissue lysate from lean and obese Ossabaw swine.

Myeloperoxidase Activity. Myeloperoxidase activity was determined in serum using previously described methods (25; 40) with modifications. Adipose tissue was homogenized in 50 mM potassium phosphate buffer (pH 6.0) with 0.5% (w/v) hexadecyltrimethylammonium bromide and freeze thawed and vortexed three times. Samples were then centrifuged for 10 minutes at 10,000 x g and the supernatants were collected. Adipose tissue supernatants and serum were mixed with o-dianisidine dihydrochloride and 0.005% (v/v) hydrogen peroxide. The oxidation of o-dianisidine was measured by the change of absorbance over a 10 minute period at 460 nm and 25°C. One unit of myeloperoxidase activity was defined as that degrading one µmol of hydrogen peroxide per min at 25°C. A molar extinction coefficient of $1.3 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ was used for determining myeloperoxidase activity.

Statistical Analysis. All statistical procedures outlined were performed using SAS 9.2 (SAS, Cary, NC). Residuals were analyzed to detect outliers and normality using the PROC Univariate procedure. Right-skewed data were log-transformed for further analysis. ANOVA was performed using the PROC MIXED procedure. Main effects for Ossabaw tissue data included lean (LFC) and obese (HFP and HFPn3) and tissue where appropriate. Fixed effects include sex, group, and termination date. Fixed effects were used in the model when $p \leq 0.15$ for the fixed effect or interaction with main effect(s). Cell culture data for

Western and NF κ B ELISA analysis were done with a cell culture lot containing an even mixture of SVC from 3 pigs from the respective breed. All other data was done on three separate SVC from each breed unless otherwise noted. SVC replicate was included in the model when $p \leq 0.15$ for the fixed effect or interaction with swine breed. Data presented are least-squares means \pm standard error. Data was considered statistically different when $p < 0.05$.

Results

To assess the ability of Ossabaw swine to accrue adipose tissue, we measured adipocyte size in SQ and VIS adipose tissue depots (**Figure 5**). Swine fed high fat diets (i.e. HFP and HFPn3) had marked enlargement adipocytes compared to the LFC fed swine SQ and VIS depots, with no difference between adipose depots. To verify comparable adipocyte size of lean Ossabaw and Yorkshire swine, we also determined adipocyte size in Yorkshire SQ adipose tissue. There was no difference in LFC Ossabaw and Yorkshire SQ adipose tissue adipocyte size (data not shown). Thus, our results demonstrate that Ossabaw swine have a marked capacity to store excess energy through adipocyte hypertrophy. Additionally, in a lean state, Ossabaw and Yorkshire swine do not differ in adipocyte size.

To compare Ossabaw and Yorkshire adipocytes *in vitro*, SVC from lean Ossabaw and Yorkshire SQ adipose tissue were proliferated and differentiated into adipocytes. Yorkshire SVC cells proliferated faster during cell expansion (**Figure 6A**). Thus, to ensure comparable cell populations, differentiation was not initiated until one day post cell confluence. During the initial stages of differentiation into adipocytes, Ossabaw SVC had increased clonal expansion compared to Yorkshire SVC (**Figure 6B**); representing an increased capacity for adipose expansion due to preadipocyte hyperplasia. Concomitant with the increased clonal

expansion, Ossabaw adipocytes also had increased enzymatic activity of G3PDH (**Figure 6C**), a marker of adipogenesis. Despite increased clonal expansion and G3PDH activity, Ossabaw adipocytes had decreased uptake of Oil Red O (**Figure 7**), and therefore decreased lipid filling compared to Yorkshire adipocytes. In addition to clonal expansion and G3PDH activity, PPAR γ 1, but not PPAR γ 2 or total PPAR γ , was increased in Ossabaw compared to Yorkshire adipocytes (**Figure 8**).

A glucose analogue, 2-NBDG, was used to test the effect of palmitate and LPS treatments on insulin sensitivity in Ossabaw and Yorkshire adipocytes (**Figure 9**). Insulin increased 2-NBDG uptake in both Ossabaw and Yorkshire adipocytes. Yorkshire adipocytes had both increased basal and insulin-stimulated 2-NBDG uptake compared to Ossabaw adipocytes. Both palmitate and LPS treatment induced insulin resistance compared vehicle treatment. However, palmitate and LPS did not differentially alter insulin resistance in either adipocyte genotype. In spite of the increased basal and insulin-stimulated 2-NBDG uptake displayed by Yorkshire adipocytes, both genotypes had a ~1.5 fold increase in 2-NBDG uptake following insulin stimulation.

Nuclear translocation of NF κ B p65 was determined in Ossabaw and Yorkshire adipocytes (**Figure 10A**). NF κ B nuclear translocation was increased by palmitate and LPS in both adipocyte genotypes. However, there was no difference in NF κ B translocation between Ossabaw and Yorkshire adipocytes. Cellular PPAR γ content was determined by western blot (**Figure 8**). Total PPAR γ , PPAR γ 1, and PPAR γ 2 content was not altered by treatment (i.e. V, P, L), however as mentioned above, Ossabaw adipocytes had increased PPAR γ 1 compared Yorkshire adipocytes.

In the media, IL-8 production by adipocytes was differentially regulated by treatment (i.e. V, P, L), with Yorkshire adipocytes producing significantly more IL-8 than Ossabaw adipocytes (**Figure 10B**). Both Ossabaw and Yorkshire adipocytes had increased IL-8 production in response to LPS, but only Yorkshire adipocytes responded to palmitate with increased IL-8 production. Yorkshire and Ossabaw adipocytes did not differ in IL-6 excretion, nor did either genotype produce IL-6 in response to palmitate (**Figure 10C**). However, like IL-8 Ossabaw adipocytes produced more IL-6 in response to LPS than Yorkshire adipocytes.

Since palmitate induced IL-8 production in Yorkshire, but not Ossabaw adipocytes, we determined whether there was any high fat diet-induced increase in serum or adipose tissue IL-8 in Ossabaw swine. There was no diet-induced increase in serum IL-8 concentrations (**Figure 11A**). Additionally, there were no diet or adipose tissue differences in IL-8 concentrations (**Figure 11B**). Although there was a significant diet*tissue interaction, with obese VIS adipose tissue having the highest IL-8 concentrations, and lean VIS adipose tissue have the lowest concentrations. Since IL-8 is a chemokine that causes neutrophil extravasation, we determined myeloperoxidase activity in serum and SQ and VIS adipose tissue. There were no differences between lean and obese pigs in serum myeloperoxidase activity (**Figure 11C**). Furthermore, myeloperoxidase activity was not detected in any adipose tissue depot of any pig.

Discussion

As evidenced by Faris *et al.* (14) and others (4; 6; 27), Ossabaw swine gain considerable weight when fed excess calories. We have demonstrated that this excessive body weight gain leads to a ~4.3 and ~8.0 fold increases in SQ and VIS adipocyte size,

respectively. Adipocyte hypertrophy in obese Ossabaw swine is much greater in magnitude compared to the ~2 to 3 fold increase in obese human (36) and rodents (25). Despite apparent obesity and adipocyte hypertrophy in SQ and VIS adipose tissue, Ossabaw swine did not display any evidence of systemic inflammation, and only a modest increase in a potentially inflammatory adipose tissue macrophage phenotype. In humans, both subcutaneous (19) and visceral (33-34) adipocyte size are positively associated with an unhealthy metabolic profile including impaired glucose homeostasis and hyperlipidemia. However, adipocyte size is not correlated with systemic inflammation measured by circulating concentrations of TNF α , IL-6, and CRP (19). In addition to no association of inflammation with adipocyte size, MHO individuals have decreased macrophage infiltration into adipose tissue (34), which likely provides protection against increased systemic inflammation. Thus, the ability to expand adipose tissue may have protected Ossabaw swine from increases in systemic inflammation, adipose tissue macrophage infiltration and consequently adipose tissue inflammation.

In a lean state, Ossabaw and Yorkshire swine have similar adipocyte size despite large differences in mature body weight. Ossabaw swine have a genetic potential to accrue and store large amounts of energy in adipose tissue due to excess caloric intake; however, genetic selection of Yorkshire swine for lean mass growth limits the ability of Yorkshire swine to store energy in adipose tissue. Potentially, phenotypical and genetic differences between Ossabaw and Yorkshire swine will translate to differences in adipocyte cell culture. Despite less lipid filling by Ossabaw SVC derived adipocytes, they have increased adipogenesis as evidenced by increased clonal expansion, higher expression of PPAR γ , and higher G3PDH activity compared to Yorkshire SVC derived adipocytes. Thus, we

hypothesized that Ossabaw adipocytes will be protected from palmitate-induced inflammation, and therefore insulin resistance, due to their ability to expand adipose tissue *in vivo*.

To our knowledge, this is the first study to evaluate the effect of palmitate in regards to NFκB activation and insulin resistance in porcine SVC derived adipocytes. Similar to 3T3 L1 adipocytes (2; 5; 11; 41), palmitate induced activation of NFκB in both Ossabaw and Yorkshire adipocytes. In spite of no differences in palmitate-induced NFκB activation between Ossabaw and Yorkshire adipocytes, IL-8 production was higher in Yorkshire adipocytes. Consistent with this, there were no obesity-linked increases in serum or adipose tissue concentrations of IL-8 in Ossabaw swine. IL-8 is a chemokine that increases neutrophil extravasation into tissue during the initial stages of the immune response, and may be involved in the initial stages of adipose tissue inflammation in obesity.

Neutrophil infiltration into adipose tissue happens in the early stages of high fat feeding in mice (13), although they are not detectable after 16 weeks. There is a paucity of information regarding neutrophil infiltration in human adipose tissue in obesity. Neutrophils were not detected in adipose tissue of severely obese women, but there was increased circulating myeloperoxidase compared to lean women (32), suggesting that while not detected in adipose tissue, neutrophil activation is a component of obesity. Ossabaw adipocytes are protected against palmitate-induced IL-8 production, and thus likely neutrophil infiltration into adipose tissue of obese swine. This is confirmed by the absence of myeloperoxidase activity in obese Ossabaw swine.

Inflammation due to palmitate can be induced by a number of factors other than NFκB, including JNK, ERK, and p38 MAPK (2; 11; 41). The lack of IL-6 production by

either Ossabaw or Yorkshire adipocytes in response to palmitate does not preclude the involvement of these factors in inducing IL-8 production, as palmitate induces NF κ B activation without subsequent IL-6 production. Induction of IL-6 by LPS in both Ossabaw and Yorkshire adipocytes demonstrate distinct differences in Tlr activation between LPS and Palmitate. Evidence from human adipocyte cell culture suggests that the majority of inflammation is from the preadipocyte fraction (9). Thus, the degree of SVC differentiation may be a determining factor in cytokine production.

Inflammation has been linked to the development of insulin resistance. Both palmitate and LPS decreased insulin-stimulated 2-NBDG uptake by adipocytes. Perhaps most interesting is the lower basal and insulin-stimulated 2-NBDG uptake by Ossabaw adipocytes, suggesting differences in capacity to uptake glucose. Faris *et al.* (14) demonstrated that lean fasted Ossabaw swine challenged with 0.25 U/kg body weight insulin have a blood glucose concentration of 60.2 mg/dL (decrease of 57.6 mg/dL) after 15 minutes, while fasted Yorkshire swine challenged with 0.1 U/kg body weight insulin have a blood glucose concentration of 32.6 mg/dL (decrease of 46.8) after 15 minutes (Faris, R.J. and M.E. Spurlock, unpublished data). This is a substantial increase in insulin sensitivity in Yorkshire swine vs. Ossabaw swine, although differences in lean body mass may affect insulin responsiveness. Nonetheless, this increase in insulin sensitivity is reflected in adipocyte cell culture.

Taken together, this data demonstrates that Ossabaw adipocytes have increased adipogenesis, and that this may be protective against inflammation as Ossabaw adipocytes did not produce IL-8 in response to palmitate. Furthermore, Ossabaw adipocytes have decreased 2-NBDG basal vs. insulin stimulated glucose uptake which is reflective of *in vivo*

evidence of insulin sensitivity in Ossabaw and Yorkshire swine. This *in vitro* evidence supports that the marked capacity for adipose expansion in Ossabaw swine to protect against high fat diet-induced inflammation, with respect to IL-8 production, but not NF κ B activation. Additionally, this data is supported by the absence of obesity-linked increases in serum and adipose tissue IL-8 concentrations.

References

1. Aguilar-Salinas CA, Garcia EG, Robles L, Riano D, Ruiz-Gomez DG, Garcia-Ulloa AC, Melgarejo MA, Zamora M, Guillen-Pineda LE, Mehta R, Canizales-Quinteros S, Tusie Luna MT, Gomez-Perez FJ. 2008. High adiponectin concentrations are associated with the metabolically healthy obese phenotype. *J Clin Endocrinol Metab* 93:4075-4079.
2. Ajuwon KM, Spurlock ME. 2005. Palmitate activates the NF-kappaB transcription factor and induces IL-6 and TNFalpha expression in 3T3-L1 adipocytes. *J Nutr* 135:1841-1846.
3. Andel M, Polak J, Kraml P, Dlouhy P, Stich V. 2009. [Chronic mild inflammation links obesity, metabolic syndrome, atherosclerosis and diabetes]. *Vnitr Lek* 55:659-665.
4. Bell LN, Lee L, Saxena R, Bemis KG, Wang M, Theodorakis JL, Vuppalanchi R, Alloosh M, Sturek M, Chalasani N. 2010. Serum proteomic analysis of diet-induced steatohepatitis and metabolic syndrome in the Ossabaw miniature swine. *Am J Physiol Gastrointest Liver Physiol* 298:G746-754.
5. Bradley RL, Fisher FF, Maratos-Flier E. 2008. Dietary fatty acids differentially regulate production of TNF-alpha and IL-10 by murine 3T3-L1 adipocytes. *Obesity (Silver Spring)* 16:938-944.
6. Bratz IN, Dick GM, Tune JD, Edwards JM, Neeb ZP, Dincer UD, Sturek M. 2008. Impaired capsaicin-induced relaxation of coronary arteries in a porcine model of the metabolic syndrome. *Am J Physiol Heart Circ Physiol* 294:H2489-2496.
7. Brochu M, Tchernof A, Dionne IJ, Sites CK, Eltabbakh GH, Sims EA, Poehlman ET. 2001. What are the physical characteristics associated with a normal metabolic profile despite a high level of obesity in postmenopausal women? *J Clin Endocrinol Metab* 86:1020-1025.
8. Burgess TA, Robich MP, Chu LM, Bianchi C, Sellke FW. 2011. Improving glucose metabolism with resveratrol in a swine model of metabolic syndrome through alteration of signaling pathways in the liver and skeletal muscle. *Arch Surg* 146:556-564.

9. Chung S, Lapoint K, Martinez K, Kennedy A, Boysen Sandberg M, McIntosh MK. 2006. Preadipocytes mediate lipopolysaccharide-induced inflammation and insulin resistance in primary cultures of newly differentiated human adipocytes. *Endocrinology* 147:5340-5351.
10. Clark BA, Alloosh M, Wenzel JW, Sturek M, Kostrominova TY. 2011. Effect of diet-induced obesity and metabolic syndrome on skeletal muscles of Ossabaw miniature swine. *Am J Physiol Endocrinol Metab* 300:E848-857.
11. Davis JE, Gabler NK, Walker-Daniels J, Spurlock ME. 2009. The c-Jun N-terminal kinase mediates the induction of oxidative stress and insulin resistance by palmitate and toll-like receptor 2 and 4 ligands in 3T3-L1 adipocytes. *Horm Metab Res* 41:523-530.
12. DiGirolamo M, Fine JB, Tagra K, Rossmanith R. 1998. Qualitative regional differences in adipose tissue growth and cellularity in male Wistar rats fed ad libitum. *Am J Physiol* 274:R1460-1467.
13. Elgazar-Carmon V, Rudich A, Hadad N, Levy R. 2008. Neutrophils transiently infiltrate intra-abdominal fat early in the course of high-fat feeding. *J Lipid Res* 49:1894-1903.
14. Faris RJ, Boddicker RL, Walker-Daniels J, Li J, Jones DE, Spurlock ME. 2012. Inflammation in response to n3 fatty acids in a porcine obesity model. *Comp Med* 62.
15. Ford ES. 2005. Prevalence of the metabolic syndrome defined by the International Diabetes Federation among adults in the U.S. *Diabetes Care* 28:2745-2749.
16. Freedman DS. 2011. Obesity - United States, 1988-2008. *MMWR Surveill Summ* 60 Suppl:73-77.
17. Gamou S, Shimizu Y, Shimizu N. 1990. *Adipocytes*. Clifton, NJ: Humana Press.
18. Hayes L, Pearce MS, Fribank MJ, Walker M, Taylor R, Unwin NC. 2010. Do obese but metabolically normal women differ in intra-abdominal fat and physical activity levels from those with the expected metabolic abnormalities? A cross-sectional study. *BMC Public Health* 10:723.
19. Hoffstedt J, Arner E, Wahrenberg H, Andersson DP, Qvist V, Lofgren P, Ryden M, Thorne A, Wiren M, Palmer M, Thorell A, Toft E, Arner P. 2010. Regional impact of adipose tissue morphology on the metabolic profile in morbid obesity. *Diabetologia* 53:2496-2503.
20. Jennings CL, Lambert EV, Collins M, Joffe Y, Levitt NS, Goedecke JH. 2008. Determinants of insulin-resistant phenotypes in normal-weight and obese Black African women. *Obesity (Silver Spring)* 16:1602-1609.
21. Karelis AD, Faraj M, Bastard JP, St-Pierre DH, Brochu M, Prud'homme D, Rabasa-Lhoret R. 2005. The metabolically healthy but obese individual presents a favorable inflammation profile. *J Clin Endocrinol Metab* 90:4145-4150.

22. Karelis AD, St-Pierre DH, Conus F, Rabasa-Lhoret R, Poehlman ET. 2004. Metabolic and body composition factors in subgroups of obesity: what do we know? *J Clin Endocrinol Metab* 89:2569-2575.
23. Kim JY, van de Wall E, Laplante M, Azzara A, Trujillo ME, Hofmann SM, Schraw T, Durand JL, Li H, Li G, Jelicks LA, Mehler MF, Hui DY, Deshaies Y, Shulman GI, Schwartz GJ, Scherer PE. 2007. Obesity-associated improvements in metabolic profile through expansion of adipose tissue. *J Clin Invest* 117:2621-2637.
24. Koster A, Stenholm S, Alley DE, Kim LJ, Simonsick EM, Kanaya AM, Visser M, Houston DK, Nicklas BJ, Tyllavsky FA, Satterfield S, Goodpaster BH, Ferrucci L, Harris TB. 2010. Body fat distribution and inflammation among obese older adults with and without metabolic syndrome. *Obesity (Silver Spring)* 18:2354-2361.
25. Kurki E, Shi J, Martonen E, Finckenberg P, Mervaala E. 2012. Distinct effects of calorie restriction on adipose tissue cytokine and angiogenesis profiles in obese and lean mice. *Nutr Metab (Lond)* 9:64.
26. Lee J, Xu Y, Lu L, Bergman B, Leitner JW, Greyson C, Draznin B, Schwartz GG. 2010. Multiple abnormalities of myocardial insulin signaling in a porcine model of diet-induced obesity. *Am J Physiol Heart Circ Physiol* 298:H310-319.
27. Lee L, Alloosh M, Saxena R, Van Alstine W, Watkins BA, Klaunig JE, Sturek M, Chalasani N. 2009. Nutritional model of steatohepatitis and metabolic syndrome in the Ossabaw miniature swine. *Hepatology* 50:56-67.
28. Martin RJ, Gobble JL, Hartsock TH, Graves HB, Ziegler JH. 1973. Characterization of an obese syndrome in the pig. *Proc Soc Exp Biol Med* 143:198-203.
29. Monteiro R, Azevedo I. 2010. Chronic inflammation in obesity and the metabolic syndrome. *Mediators Inflamm* 2010.
30. Neeb ZP, Edwards JM, Alloosh M, Long X, Mokelke EA, Sturek M. 2010. Metabolic syndrome and coronary artery disease in Ossabaw compared with Yucatan swine. *Comp Med* 60:300-315.
31. Nguyen NT, Nguyen XM, Lane J, Wang P. 2011. Relationship between obesity and diabetes in a US adult population: findings from the National Health and Nutrition Examination Survey, 1999-2006. *Obes Surg* 21:351-355.
32. Nijhuis J, Rensen SS, Slaats Y, van Dielen FM, Buurman WA, Greve JW. 2009. Neutrophil activation in morbid obesity, chronic activation of acute inflammation. *Obesity (Silver Spring)* 17:2014-2018.
33. O'Connell J, Lynch L, Cawood TJ, Kwasnik A, Nolan N, Geoghegan J, McCormick A, O'Farrelly C, O'Shea D. 2010. The relationship of omental and subcutaneous adipocyte size to metabolic disease in severe obesity. *PLoS One* 5:e9997.
34. O'Connell J, Lynch L, Hogan A, Cawood TJ, O'Shea D. 2011. Preadipocyte factor-1 is associated with metabolic profile in severe obesity. *J Clin Endocrinol Metab* 96:E680-684.

35. Robich MP, Osipov RM, Chu LM, Han Y, Feng J, Nezafat R, Clements RT, Manning WJ, Sellke FW. 2011. Resveratrol modifies risk factors for coronary artery disease in swine with metabolic syndrome and myocardial ischemia. *Eur J Pharmacol* 664:45-53.
36. Salans LB, Cushman SW, Weismann RE. 1973. Studies of human adipose tissue. Adipose cell size and number in nonobese and obese patients. *J Clin Invest* 52:929-941.
37. Shin MJ, Hyun YJ, Kim OY, Kim JY, Jang Y, Lee JH. 2006. Weight loss effect on inflammation and LDL oxidation in metabolically healthy but obese (MHO) individuals: low inflammation and LDL oxidation in MHO women. *Int J Obes (Lond)* 30:1529-1534.
38. Sims EA. 2001. Are there persons who are obese, but metabolically healthy? *Metabolism* 50:1499-1504.
39. Stefan N, Kantartzis K, Machann J, Schick F, Thamer C, Rittig K, Balletshofer B, Machicao F, Fritsche A, Haring HU. 2008. Identification and characterization of metabolically benign obesity in humans. *Arch Intern Med* 168:1609-1616.
40. Suzuki K, Ota H, Sasagawa S, Sakatani T, Fujikura T. 1983. Assay method for myeloperoxidase in human polymorphonuclear leukocytes. *Anal Biochem* 132:345-352.
41. Wang SL, Li Y, Wen Y, Chen YF, Na LX, Li ST, Sun CH. 2009. Curcumin, a potential inhibitor of up-regulation of TNF-alpha and IL-6 induced by palmitate in 3T3-L1 adipocytes through NF-kappaB and JNK pathway. *Biomed Environ Sci* 22:32-39.
42. Wangsness PJ, Martin RJ, Gahagan JH. 1977. Insulin and growth hormone in lean and obese pigs. *Am J Physiol* 233:E104-108.
43. Wangsness PJ, Martin RJ, Gatchel BB. 1980. Insulin induced growth hormone response in fast-growing, lean and in slow-growing, obese pigs. *Growth* 44:318-326.
44. Weiss R, Taksali SE, Dufour S, Yeckel CW, Papademetris X, Cline G, Tamborlane WV, Dziura J, Shulman GI, Caprio S. 2005. The "obese insulin-sensitive" adolescent: importance of adiponectin and lipid partitioning. *J Clin Endocrinol Metab* 90:3731-3737.
45. Yki-Jarvinen H. 2005. Fat in the liver and insulin resistance. *Ann Med* 37:347-356.

Figures




Figure 5. Comparison of SQ and VIS adipocyte size in lean and obese Ossabaw Swine.

(A) Adipocyte area of SQ and VIS adipose tissue sections from lean (LFC) and obese (HFP and HFPn3) Ossabaw swine and examples of (B) lean and (C) obese Ossabaw swine adipose tissue sections. Data presented is Least-Square Means \pm standard error. Main Effects: (A) Diet $p < 0.0001$, Tissue $P = 0.31$, Diet*Tissue $p = 0.0315$.




Figure 6. *Ossabaw adipocytes have increased clonal expansion and G3PDH activity compared to Yorkshire adipocytes.*

(A) BrDU incorporation during growth phase and (B) differentiation phase of Ossabaw and Yorkshire SVC. (C) G3PDH activity of Ossabaw and Yorkshire adipocytes 10 days post differentiation. Data presented is Least-Square Means \pm standard error. Means with different letters represent a difference of $p < 0.05$. Main Effects: (A) Genotype $p < 0.0001$, Day $p < 0.0001$, Genotype*Day $p = 0.0048$; (B) Genotype $p < 0.0001$, Day $p < 0.0001$, Genotype*Day $p < 0.0001$; (C) Genotype $p = 0.0104$. Each experiment was replicated three times with an $n = 4$ for each experiment.




Figure 7. *Yorkshire adipocytes have more lipid filling than Ossabaw adipocytes.*

(A) Oil Red O extracted from Ossabaw and Yorkshire adipocytes 10 days post-differentiation. Representative (B) Ossabaw and (C) Yorkshire Oil Red O images. Data presented is Least-Square Means \pm standard error. Means with different letters represent a difference of $p < 0.05$. Main Effect: (A) Genotype $p < 0.0001$. Each experiment was replicated three times with an $n=6$ for each experiment.




Figure 8. *Ossabaw adipocytes express more cellular PPARγ1 than Yorkshire adipocytes.*

Cellular PPARγ1, PPARγ2, total PPARγ, and representative blot in Ossabaw and Yorkshire adipocytes treated with Vehicle, Palmitate, and LPS. There was no difference by treatment. Data presented is Least-Square Means \pm standard error for Ossabaw and Yorkshire adipocytes. * Represents differences ($p < 0.05$) between Ossabaw and Yorkshire adipocytes. Main Effects: (PPARγ1) Genotype $p = 0.0151$, Treatment $p = 0.33$, Genotype*Treatment $p = 0.40$; (PPARγ2) Genotype $p = 0.13$, Treatment $p = 0.51$, Genotype*Treatment $p = 0.81$; (PPARγ) Genotype $p = 0.13$, Treatment $p = 0.29$, Genotype*Treatment $p = 0.43$. PPARγ assay was performed on an evenly mixed SVC lot of three pigs with an $n = 4$.




Figure 9. *Ossabaw adipocytes have decreased insulin sensitivity compared to Yorkshire adipocytes, but do not respond differently to palmitate or LPS treatment.*

Insulin stimulated 2-NBDG uptake in Ossabaw and Yorkshire adipocytes treated with Vehicle (V), Palmitate (P), and LPS (L). Data presented is Least-Square Means \pm standard error. Main Effects: Treatment $p < 0.0001$; Genotype $p < 0.0001$, Treatment*Genotype $p = 0.33$. Each experiment was replicated three times with an $n = 4$ for each experiment.




Figure 10. *Palmitate and LPS differentially regulate inflammation in Ossabaw and Yorkshire adipocytes.*

(A) NFκB p65 nuclear translocation, (B) media IL-8, and (C) media IL-6 in Ossabaw and Yorkshire adipocytes treated with Vehicle (V), Palmitate (P), and LPS (L). Data presented is Least-Square Means \pm standard error. Different letter represent a difference of $p < 0.05$. Main Effects: (A) Treatment $p < 0.0001$, Genotype $p = 0.86$, Treatment*Genotype $p = 0.35$; (B) Treatment $p < 0.0001$, Genotype $p = 0.0037$, Treatment*Genotype $p < 0.0001$; (C) Treatment $p < 0.0001$, Genotype $p = 0.0396$, Treatment*Genotype $p = 0.0143$. (A) NFκB assay was performed on an evenly mixed SVC lot of three pigs with an $n = 5$. (B) IL-8 and (C) IL-6 assays were replicated 3 times with an $n = 4$ for each experiment.




Figure 11. *Ossabaw swine do not have obesity-linked increases in IL-8 or myeloperoxidase activity.*

(A) Serum IL-8, (B) SQ and VIS adipose tissue IL-8 and (C) serum myeloperoxidase activity of lean and obese Ossabaw swine. Data presented is Least-Square Means \pm standard error. Different letter represent a difference of $p < 0.05$. Main Effects: (A) Diet $p = 0.63$; (B) Diet $p = 0.90$, Tissue $p = 0.21$, Diet*Tissue $p = 0.0004$; (C) Diet $p = 0.65$.

CHAPTER 5: ADIPOSE TISSUE DEPOT DIFFERENCES IN MICROARRAY EXPRESSION IN OSSABAW SWINE

Abstract

Obesity is characterized by the increase in adipose tissue mass which can occur in multiple adipose tissue depots. In particular, visceral adiposity is associated with an increased risk for the development of type 2 diabetes and cardiovascular disease. We determined the global gene expression of subcutaneous and visceral adipose tissue by Affymetrix microarray analysis under conditions of restricted (n=7) or *ad libitum* (n=7) feed intake for 38 weeks in Ossabaw swine. No genes were determined to be differentially expressed by dietary treatment, however, 1494 were determined to be differentially expressed ($Q < 0.05$) between subcutaneous and visceral adipose tissue, of which 777 were annotated. Gene ontology and KEGG Pathway analysis were performed using DAVID. The gene ontology cellular component category extracellular region was determined to be over-represented with 36 genes by depot. The over-represented KEGG pathways identified included arachidonic acid metabolism and chemokine signaling pathway by depot. These genes and pathways identify distinct differences in subcutaneous and visceral adipose tissue that may be important in obesity.

Introduction

Obesity, by definition, is the increase in adipose tissue mass which can occur in multiple anatomically distinct adipose tissue depots (i.e. subcutaneous, retroperitoneal, mesenteric, omental, etc.). In addition to adipose tissue depots, obesity may lead to ectopic lipid deposition in other tissues, particularly the liver (11), muscle (36), and pancreas (38). Waist circumference, which is reflective of abdominal or visceral adiposity and is a component of the metabolic syndrome (1), has been increasing in both men and women in the United States (29). Visceral adiposity increases the risk of development of the metabolic syndrome (27; 31), and thus increases the risk for development of type 2 diabetes (16) and cardiovascular disease (18; 31).

Anatomical location of the adipose tissue depot is a determinant in the functional properties of the tissue. Enzymes involved in fatty acid metabolism are differentially regulated between adipose tissue depots. Visceral adipocytes have increased acyl-Co-A synthetase (2) and diacylglycerol acyltransferase (DGAT) (2; 22) activity compared with subcutaneous adipocytes in overweight individuals, but DGAT is not increased in obese individual's visceral adipocytes (22). In addition to increased enzyme activities, visceral adipose tissue has increased inflammatory gene expression. Compared to subcutaneous adipose tissue, visceral adipose tissue has increased expression of 12/15 lipoxgenase, interleukin (IL) 6 , IL-12 p35, CXCL10 (10) , CCL2 (5), and IL-8 (4). Additionally, macrophage secreted factors increase IL-6 production more in visceral adipocytes than subcutaneous adipocytes (35). The increased inflammatory profile of visceral adipocytes is important because it can contribute directly to increased liver production of C-reactive protein (CRP) (3), and further the chronic low-grade inflammation associated with obesity.

The need for alternative comparative models to investigate differences in adipose depot gene expression under lean and obese conditions is evident. Murine models have been pivotal to investigating the etiology of obesity and the roles of specific genes. However, human and mice differ on a number of factors relating to studying adipose tissue and obesity. Specifically, CRP is not highly expressed nor is it an acute phase protein in mice (39), human macrophages do not produce nitric oxide (32), and gonadal fat is not a major adipose tissue depot in humans. Swine are a viable alternative due to their comparative physiology and anatomy. Ossabaw swine are particular interest because of their ability to accumulate fat mass and their previous use in studying obesity and its co-morbidities (12; 15; 28). Thus, the objective of this research was to determine global gene expression in subcutaneous and visceral adipose tissue in lean and obese Ossabaw swine.

Methods

Animals. All animal procedures were conducted at Iowa State University and were approved by the Iowa State University Institutional Animal Care and Use Committee. Fourteen sexually mature male and female Ossabaw swine (*Sus scrofa*) (average initial body weight 46.7 kg) were housed in individual pens in a temperature controlled environment with a 12 hour light cycle. Swine were obtained from the Comparative Medicine Program at the Indiana University School of Medicine and Purdue University from a herd that tested negative for *Brucella* spp. and psuedorabies, and showed no clinical evidence of communicable disease upon arrival or during the study. Swine were fed a diet formulated to meet minimum requirements for mature swine (33) for 38 weeks. Dietary treatments were *ad libitum* feed (n=7; 4 male, 3 female) or restricted feed (n=7, 3 male, 4 female). After 38 weeks of dietary treatment, animals were fasted 12 hours before blood collection via jugular

venipuncture. Thereafter, animals were anesthetized with telazol:ketamine:xylazine (1:1:1) and then were euthanized by exsanguination. Subcutaneous (SQ) adipose tissue from back fat above the 10th rib and visceral adipose tissue (VIS) surrounding the intestines were collected and snap frozen in liquid nitrogen.

Serum analysis. Cholesterol, glucose, triglycerides, and non-esterified fatty acids (NEFA) were analyzed using Chol Slides (Ortho Clinical Diagnostics, Rochester, NY), Glu slides (Ortho Clinical Diagnostics), Trig Slides (Ortho Clinical Diagnostics), and NEFA-HR (Wako Diagnostics, Richmond, VA), respectively, on a Vitros 5.1 Chemistry Analyzer (Ortho Clinical Diagnostics). Serum insulin concentrations were measured by EIA (ALPCO, Salem, NH). The HOMA-IR was calculated using fasting serum insulin ($\mu\text{U/mL}$) x fasting blood glucose (mg/dL) divided by 405 (30). Serum endotoxin was determined using Chromogenic LAL Assay (Lonza, Anaheim, CA).

RNA Isolation. RNA was isolated from SQ and VIS adipose tissue by Guanidinium Thiocyanate-Phenol-Chloroform extraction as previously described (7). Quality and quantity of RNA was analyzed using an Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA). Microarray analysis was performed at the Iowa State University GeneChip Facility (Ames, IA) using Affymetrix Porcine GeneChip Genome Array according to manufacturer's directions (Affymetrix, Santa Clara, CA).

Microarray analysis. Transcript levels were normalized to the chip median and log transformed. For each probeset, the generalized linear model $Y_{ij} = \mu + B_1 T_i + \varepsilon_{ij}$ was fit. In each ANOVA, Y_{ij} is the log normalized transcript level for the i^{th} treatment and the j^{th} replicate, μ is the overall mean expression for the feature and T_i represents the i^{th} treatment (Treatment, Depot, Sex, Treatment*Depot, Treatment*Sex, Depot*Sex,

Treatment*Depot*Sex). An F test of the effect of treatment for each probeset was conducted as the ratio of the mean squares for treatment over the mean squares for error, and the p value for the test of the null hypothesis (i.e., mean expression not different among the treatments) was calculated. We examined the model for conformation to the assumption of normality of the residuals testing the null hypothesis that the residuals for each gene were normally distributed using the Shapiro-Wilkes Test. For each p-value, a Q-value was determined to control for FDR. All analyses were performed in SAS 9.2 (SAS Institute, Cary, NC). If the test of the null hypothesis of difference across treatments was rejected, and we had no evidence for departure from normality of the residuals, we declared the gene differentially expressed across treatments. Microarray data was annotated with gene title, gene symbol, and Entrez gene ID (9) (Hu, Z., J. Reecy, and C. Tuggle. Unpublished data). DAVID (23-24) was used to determine over-represented (EASE Score $p < 0.05$) biological process, cellular component, and molecular functional categories using the GO FAT databases. Additionally, over-represented KEGG pathways were also determined.

Statistical analysis. Statistical procedures for serum analysis was performed using SAS 9.2 (SAS, Cary, NC). Residuals were analyzed to detect outliers and normality using the PROC Univariate procedure. Right-skewed data were log-transformed for further analysis. ANOVA was performed using the PROC MIXED procedure. The main effect was dietary treatment. Sex was considered a fixed effect and was included in the model when $p \leq 0.15$ for the fixed effect or interaction with dietary treatment.

Results

After 38 weeks of dietary treatment *ad libitum* fed Ossabaw swine became obese weighing an average of 118.8 kg compared to the restricted fed Ossabaw swine weighing an

average of 59.8 kg. At the end of the study, obese Ossabaw swine were metabolically healthy (**Table 5**), with was no differences in circulating serum concentrations of insulin, glucose, NEFA, or endotoxin. Nor was there a difference in HOMA-IR. Both serum cholesterol and triglycerides were increased by dietary treatment, although the increase likely was not pathological.

To determine the effect of dietary treatment, adipose tissue depot, sex, and their interactions, microarray analysis was performed on SQ and VIS adipose tissue. At total of 19,540 genes were determined to be expressed and included in further analysis. Of these, a total of 9617 genes were successfully annotated. After controlling for FDR, only 7 genes by sex and 1494 genes by depot were determined differentially expressed ($Q < 0.05$). Only one gene (thymosin beta 4, X linked) out of 7 was annotated for sex effect. For the depot effect, 777 out of 1494 differentially expressed genes ($Q < 0.05$) were annotated. At a cutoff of $Q < 0.01$, 95 out of 185 genes were annotated. This data is presented in **Table 6**, with the Affy ID, gene title, Q-value, and fold change. Of these genes, a total of 38 genes were up-regulated in VIS adipose tissue, relative to SQ adipose tissue, while 57 were down regulated.

To determine over-represented gene categories and pathways, the 777 annotated differentially expressed genes ($Q < 0.05$) were analyzed by DAVID using the Entrez Gene ID for gene ontology molecular function, cellular component, and biological process categories (GOFAT) and the KEGG pathway database. The Entrez Gene IDs for all annotated genes were used as the background list. In all, only one cellular component (extracellular region) category and two KEGG pathways (arachidonic acid metabolism and chemokine signaling pathway) were determined to be over-represented (EASE Score < 0.05) by differently expressed genes by depot. Genes expressed in extracellular region, arachidonic acid

metabolism, and chemokine signaling pathway categories are listed in **Tables 7-9**, respectively, and will be discussed further in the next section.

Discussion

Despite the duration of the study and body weight gain by the *ad libitum* fed Ossabaw swine, they did not display a perturbed metabolic profile. Nor did they have any differentially expressed genes by dietary treatment after controlling for FDR. Dietary factors may be casual to the development of an un-healthy metabolic profile, and the absence of high dietary fat may have precluded the development of altered glucose homeostasis, hyperlipidemia, and altered adipose tissue gene expression. As evidenced by others (12; 28), the inclusion of high dietary fat and supra-physiological levels of cholesterol induces hyperlipidemia in Ossabaw swine. Additionally, high dietary fat in itself is enough to induce insulin resistance in Ossabaw swine (15). Nonetheless, distinct differences between SQ and VIS depots were evident by microarray analysis.

The gene ontology cellular component extracellular region refers to gene products that are secreted outside the cell. Thirty-six genes were over-represented in this category (**Table 7**) from the significant differentially expressed genes by depot. Four of these genes are involved in the insulin-like growth factor (IGF) axis: IGF1, IGFBP3, IGFBP5, and IGFBP6. There is a relative paucity of information relating to IGF-1 and its binding protein in regards to different adipose tissue depots. IGF-1 is important in both the proliferation of preadipocytes and differentiation into adipocytes (20; 37). However, IGF-1-induced PKB signaling is impaired in visceral adipocytes compared to subcutaneous adipocytes (8). IGF-1 and IGFBP-3 are decreased in response to increased body fat, especially in women (19). Additionally, increasing IGF-1 concentrations via growth hormone supplementation

decreases visceral adiposity (17; 21; 34). Whether this is because of increasing IGF-1 or directly to growth hormone is unclear. Potentially, lower concentrations of IGF-1 and IGFBP in visceral adipose tissue result in decreased proliferation and cell survival and thus, contribute to adipose tissue dysfunction.

In addition to IGF proteins, three chemokines were also identified as over-represented genes in the extracellular region gene ontology cellular component category (**Table 7**). These chemokines are CCL2, CXCL2, and CCL21. CCL2 and CXCL2 were down-regulated in visceral adipose tissue compared to subcutaneous whereas CCL21 was up-regulated. Chemokine signaling pathway was also identified as a KEGG pathway containing over-represented genes (**Table 9**), including CCL2, CXCL2, CCL21, and CCR3. CCL2 is increased in obesity (5), and its importance in macrophage infiltration into adipose tissue is well established (26). In humans, CCL2 is produced at a higher rate in visceral than subcutaneous adipose tissue, however, after adjusting for tissue macrophages this difference disappeared (5). Based upon the lack of metabolic and microarray differences by dietary treatment, it is likely that the decreased CCL2 expression in visceral adipose tissue is due to lack of pathological consequences from diet induced obesity and only reflects strictly depot differences. Other than increased expression of CCR3 in subcutaneous adipose tissue in obesity (25), literature pertaining to CCR3, CXCL2, and CCL21 pertaining to obesity and adipose tissue depots is absent.

In addition to chemokine signaling pathway, seven genes were identified in the arachidonic acid metabolism KEGG Pathway as containing over-represented genes by depot in our microarray analysis (**Table 8**). Products of arachidonic acid metabolism by 12/15 lipoxygenase induce inflammation and insulin resistance in 3T3 L1 adipocytes (6). Like

humans, our data demonstrates that arachidonate 12/15 lipoxygenase is more highly expressed in visceral than subcutaneous adipose tissue (10). Literature relating to obesity and adipose tissue is limited for the other gene indicated in arachidonic acid metabolism KEGG pathway. One gene, prostaglandin G/H synthase-2 (i.e. COX-2) has been implicated in obesity in rodent models (13). In humans, COX-2 expression is higher in epicardial adipose tissue than subcutaneous, but lower than that of visceral adipose tissue (14).

While global gene expression analysis did not identify genes differentially regulated by dietary treatment, it did identify differences in mRNA abundance of genes related to excreted proteins, arachidonic acid metabolism, and chemokine signaling pathway between SQ and VIS adipose tissue. These genes may be important with respect to obesity and adipose tissue inflammation and dysfunction. Further investigation into these genes in VIS and SQ adipose tissue with a high fat dietary treatment is warranted.

References

1. Alberti KG, Zimmet P, Shaw J. 2005. The metabolic syndrome--a new worldwide definition. *Lancet* 366:1059-1062.
2. Ali AH, Koutsari C, Mundi M, Stegall MD, Heimbach JK, Taler SJ, Nygren J, Thorell A, Bogachus LD, Turcotte LP, Bernlohr D, Jensen MD. 2011. Free fatty acid storage in human visceral and subcutaneous adipose tissue: role of adipocyte proteins. *Diabetes* 60:2300-2307.
3. Bastard JP, Jardel C, Delattre J, Hainque B, Bruckert E, Oberlin F. 1999. Evidence for a link between adipose tissue interleukin-6 content and serum C-reactive protein concentrations in obese subjects. *Circulation* 99:2221-2222.
4. Bruun JM, Lihn AS, Madan AK, Pedersen SB, Schiott KM, Fain JN, Richelsen B. 2004. Higher production of IL-8 in visceral vs. subcutaneous adipose tissue. Implication of nonadipose cells in adipose tissue. *Am J Physiol Endocrinol Metab* 286:E8-13.
5. Bruun JM, Lihn AS, Pedersen SB, Richelsen B. 2005. Monocyte chemoattractant protein-1 release is higher in visceral than subcutaneous human adipose tissue (AT): implication of macrophages resident in the AT. *J Clin Endocrinol Metab* 90:2282-2289.

6. Chakrabarti SK, Cole BK, Wen Y, Keller SR, Nadler JL. 2009. 12/15-lipoxygenase products induce inflammation and impair insulin signaling in 3T3-L1 adipocytes. *Obesity (Silver Spring)* 17:1657-1663.
7. Chomczynski P, Sacchi N. 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162:156-159.
8. Cleveland-Donovan K, Maile LA, Tsiaras WG, Tchkonina T, Kirkland JL, Boney CM. 2010. IGF-I activation of the AKT pathway is impaired in visceral but not subcutaneous preadipocytes from obese subjects. *Endocrinology* 151:3752-3763.
9. Couture O, Callenberg K, Koul N, Pandit S, Younes R, Hu ZL, Dekkers J, Reecy J, Honavar V, Tuggle C. 2009. ANEXdb: an integrated animal ANnotation and microarray EXpression database. *Mamm Genome* 20:768-777.
10. Dobrian AD, Lieb DC, Ma Q, Lindsay JW, Cole BK, Ma K, Chakrabarti SK, Kuhn NS, Wohlgemuth SD, Fontana M, Nadler JL. 2010. Differential expression and localization of 12/15 lipoxygenases in adipose tissue in human obese subjects. *Biochem Biophys Res Commun* 403:485-490.
11. Donnelly KL, Smith CI, Schwarzenberg SJ, Jessurun J, Boldt MD, Parks EJ. 2005. Sources of fatty acids stored in liver and secreted via lipoproteins in patients with nonalcoholic fatty liver disease. *J Clin Invest* 115:1343-1351.
12. Dyson MC, Alloosh M, Vuchetich JP, Mokelke EA, Sturek M. 2006. Components of metabolic syndrome and coronary artery disease in female Ossabaw swine fed excess atherogenic diet. *Comp Med* 56:35-45.
13. Fain JN, Ballou LR, Bahouth SW. 2001. Obesity is induced in mice heterozygous for cyclooxygenase-2. *Prostaglandins Other Lipid Mediat* 65:199-209.
14. Fain JN, Sacks HS, Bahouth SW, Tichansky DS, Madan AK, Cheema PS. 2010. Human epicardial adipokine messenger RNAs: comparisons of their expression in substernal, subcutaneous, and omental fat. *Metabolism* 59:1379-1386.
15. Faris RJ, Boddicker RL, Walker-Daniels J, Li J, Jones DE, Spurlock ME. 2012. Inflammation in response to n3 fatty acids in a porcine obesity model. *Comp Med* 62.
16. Ford ES, Schulze MB, Pischon T, Bergmann MM, Joost HG, Boeing H. 2008. Metabolic syndrome and risk of incident diabetes: findings from the European Prospective Investigation into Cancer and Nutrition-Potsdam Study. *Cardiovasc Diabetol* 7:35.
17. Franco C, Brandberg J, Lonn L, Andersson B, Bengtsson BA, Johannsson G. 2005. Growth hormone treatment reduces abdominal visceral fat in postmenopausal women with abdominal obesity: a 12-month placebo-controlled trial. *J Clin Endocrinol Metab* 90:1466-1474.
18. Galassi A, Reynolds K, He J. 2006. Metabolic syndrome and risk of cardiovascular disease: a meta-analysis. *Am J Med* 119:812-819.
19. Gomez JM, Maravall FJ, Gomez N, Navarro MA, Casamitjana R, Soler J. 2004. The IGF-I system component concentrations that decrease with ageing are lower in

- obesity in relationship to body mass index and body fat. *Growth Horm IGF Res* 14:91-96.
20. Grohmann M, Sabin M, Holly J, Shield J, Crowne E, Stewart C. 2005. Characterization of differentiated subcutaneous and visceral adipose tissue from children: the influences of TNF-alpha and IGF-I. *J Lipid Res* 46:93-103.
 21. Hong JW, Park JK, Lim CY, Kim SW, Chung YS, Lee EJ. 2011. A weekly administered sustained-release growth hormone reduces visceral fat and waist circumference in abdominal obesity. *Horm Metab Res* 43:956-961.
 22. Hou XG, Moser S, Sarr MG, Thompson GB, Que FG, Jensen MD. 2009. Visceral and subcutaneous adipose tissue diacylglycerol acyltransferase activity in humans. *Obesity (Silver Spring)* 17:1129-1134.
 23. Huang da W, Sherman BT, Lempicki RA. 2009. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 37:1-13.
 24. Huang da W, Sherman BT, Lempicki RA. 2009. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4:44-57.
 25. Huber J, Kiefer FW, Zeyda M, Ludvik B, Silberhumer GR, Prager G, Zlabinger GJ, Stulnig TM. 2008. CC chemokine and CC chemokine receptor profiles in visceral and subcutaneous adipose tissue are altered in human obesity. *J Clin Endocrinol Metab* 93:3215-3221.
 26. Kanda H, Tateya S, Tamori Y, Kotani K, Hiasa K, Kitazawa R, Kitazawa S, Miyachi H, Maeda S, Egashira K, Kasuga M. 2006. MCP-1 contributes to macrophage infiltration into adipose tissue, insulin resistance, and hepatic steatosis in obesity. *J Clin Invest* 116:1494-1505.
 27. Kim LJ, Nalls MA, Eiriksdottir G, Sigurdsson S, Launer LJ, Koster A, Chaves PH, Jonsdottir B, Garcia M, Gudnason V, Harris TB. 2011. Associations of visceral and liver fat with the metabolic syndrome across the spectrum of obesity: the AGES-Reykjavik study. *Obesity (Silver Spring)* 19:1265-1271.
 28. Lee L, Alloosh M, Saxena R, Van Alstine W, Watkins BA, Klaunig JE, Sturek M, Chalasani N. 2009. Nutritional model of steatohepatitis and metabolic syndrome in the Ossabaw miniature swine. *Hepatology* 50:56-67.
 29. Li C, Ford ES, McGuire LC, Mokdad AH. 2007. Increasing trends in waist circumference and abdominal obesity among US adults. *Obesity (Silver Spring)* 15:216-224.
 30. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. 1985. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412-419.
 31. Meigs JB, Wilson PW, Fox CS, Vasan RS, Nathan DM, Sullivan LM, D'Agostino RB. 2006. Body mass index, metabolic syndrome, and risk of type 2 diabetes or cardiovascular disease. *J Clin Endocrinol Metab* 91:2906-2912.

32. Mestas J, Hughes CC. 2004. Of mice and not men: differences between mouse and human immunology. *J Immunol* 172:2731-2738.
33. National Research Council (U.S.). Subcommittee on Swine Nutrition. 1998. *Nutrient requirements of swine*. Washington, D.C.: National Academy Press.
34. Pasarica M, Zachwieja JJ, Dejonge L, Redman S, Smith SR. 2007. Effect of growth hormone on body composition and visceral adiposity in middle-aged men with visceral obesity. *J Clin Endocrinol Metab* 92:4265-4270.
35. Permana PA, Zhang W, Wabitsch M, Fischer-Posovszky P, Duckworth WC, Reaven PD. 2009. Pioglitazone reduces inflammatory responses of human adipocytes to factors secreted by monocytes/macrophages. *Am J Physiol Endocrinol Metab* 296:E1076-1084.
36. Phillips DI, Caddy S, Ilic V, Fielding BA, Frayn KN, Borthwick AC, Taylor R. 1996. Intramuscular triglyceride and muscle insulin sensitivity: evidence for a relationship in nondiabetic subjects. *Metabolism* 45:947-950.
37. Scavo LM, Karas M, Murray M, Leroith D. 2004. Insulin-like growth factor-I stimulates both cell growth and lipogenesis during differentiation of human mesenchymal stem cells into adipocytes. *J Clin Endocrinol Metab* 89:3543-3553.
38. Shimabukuro M, Zhou YT, Levi M, Unger RH. 1998. Fatty acid-induced beta cell apoptosis: a link between obesity and diabetes. *Proc Natl Acad Sci U S A* 95:2498-2502.
39. Szalai AJ, McCrory MA. 2002. Varied biologic functions of C-reactive protein: lessons learned from transgenic mice. *Immunol Res* 26:279-287.

Tables

Table 5. *Metabolic profile of Ossabaw swine.*

Parameter	Restricted	Obese	p value
Insulin (pg/mL)	119.7±25.8	130.1±25.8	0.78
Glucose (mg/dL)	93.6±16.2	106.9±16.2	0.57
HOMA-IR	0.57±0.15	0.75±0.15	0.42
Cholesterol (mg/dL)	63.4±5.6	81.0±5.6	0.0463
Triglycerides (mg/dL)	21.6±2.1	37.5±2.1	0.0005
NEFA (mM)	0.28±0.06	0.43±0.06	0.13
Endotoxin (ng/mL)	14.9±1.3	14.7±1.3	0.91

Table 6. *Differentially expressed genes by depot ($Q < 0.01$).*

Affy ID	Rank^a	GENE TITLE	Q-value	Fold Change^b
Ssc.24864.1.A1_at	1	hypothetical LOC100515948	0.0007	13.40
Ssc.8623.1.A1_at	4	Sec23 homolog A (S. cerevisiae)	0.0014	-5.08
Ssc.5713.1.S1_at	7	gelatinase A	0.0015	-50.51
Ssc.3450.1.S1_at	8	homeobox protein Hox-D8-like	0.0029	-4.87
Ssc.12936.1.A1_at	10	keratocan-like	0.0033	-5.70
Ssc.18608.1.S1_at	15	cellular retinoic acid binding protein 1	0.0039	1.60
Ssc.15923.1.S1_at	18	fibroblast growth factor 7	0.0039	-0.25
Ssc.25680.1.S1_at	19	mutS homolog 3 (E. coli)	0.0040	0.20
Ssc.21925.1.S1_at	22	complement C1q tumor necrosis factor-related protein 5-like	0.0040	-3.53
Ssc.2445.1.S1_at	23	microfibrillar-associated protein 2-like	0.0040	-5.72
Ssc.2464.1.S1_at	25	Stanniocalcin 1	0.0040	-4.30
Ssc.21549.1.A1_at	28	meningioma expressed antigen 5 (hyaluronidase)	0.0040	0.89
Ssc.15266.1.S1_at	32	15-hydroxyprostaglandin dehydrogenase [NAD ⁺]-like	0.0042	1.80
Ssc.6327.1.S1_at	33	keratin, type I cytoskeletal 19-like	0.0042	5.95
Ssc.21125.1.S1_at	35	oral-facial-digital syndrome 1	0.0045	-1.02
Ssc.30175.1.A1_at	36	carboxypeptidase X (M14 family), member 1	0.0045	-8.90
Ssc.16418.2.S1_at	37 ^c	integrin beta-like protein 1-like	0.0052	-3.94
Ssc.16589.1.S1_at	41	collagen, type VI, alpha 3	0.0053	-48.52
Ssc.8573.1.A1_at	43	apolipoprotein D	0.0053	-40.40
Ssc.30.1.S1_at	46	proteasome (prosome, macropain) subunit, alpha type, 4	0.0053	-7.89
Ssc.19688.1.S1_at	47	cholinergic receptor, nicotinic, alpha 1 (muscle)	0.0053	-23.30
Ssc.12795.1.S1_at	48	three prime repair exonuclease 1-like	0.0055	-1.10
Ssc.24330.1.S1_at	49	cytochrome P450, family 7, subfamily B, polypeptide 1	0.0056	2.89
Ssc.4044.1.A1_at	52	hypothetical LOC100512560	0.0059	2.62
Ssc.26893.1.A1_at	53	hemicentin-1-like	0.0059	2.64
Ssc.26617.1.A1_at	54	paired mesoderm homeobox protein 1-like	0.0059	-5.25

Table 6. *Differentially expressed genes by depot ($Q < 0.01$) (continued).*

Ssc.5396.1.S1_at	59	Similar to transcription factor 21	0.0062	3.13
Ssc.24864.2.S1_at	60	hypothetical LOC100515948	0.0062	2.08
Ssc.9754.1.A1_at	62	Ankyrin repeat and SOCS box-containing 6	0.0064	-0.44
Ssc.24250.1.S1_at	63	cellular retinoic acid binding protein 2	0.0065	-11.22
Ssc.18278.2.A1_at	66	nebulin-related anchoring protein	0.0067	-0.45
Ssc.6058.1.S1_at	67	dual specificity phosphatase 1	0.0067	-34.67
Ssc.6817.1.A1_a_at	73	tumor suppressor candidate 3-like	0.0068	1.93
Ssc.2747.1.S1_at	74	homeobox protein Hox-B5-like	0.0068	1.74
Ssc.1600.3.S1_at	78	ribosomal protein S6 modification-like protein B-like	0.0068	1.21
Ssc.226.1.S1_at	83	high-mobility group box 2	0.0068	1.61
Ssc.26825.1.S1_at	85	zinc finger protein 330-like	0.0068	1.00
Ssc.14851.1.S1_at	86	amyloid beta (A4) precursor-like protein 2	0.0068	0.87
Ssc.11952.1.S1_at	87 ^c	transmembrane protein 117-like	0.0068	-1.23
Ssc.17345.1.S1_at	88	angiopoietin-like 4	0.0068	-2.56
Ssc.19323.1.S1_at	89	protein NOXP20-like	0.0070	-6.46
Ssc.24429.1.A1_at	90	sperm-specific antigen 2-like	0.0070	-0.20
Ssc.1310.1.S1_at	91	prostaglandin E synthase	0.0070	-4.09
Ssc.19136.1.S1_at	93	latent-transforming growth factor beta-binding protein 2-like	0.0071	1.73
Ssc.11369.2.A1_at	97	Der1-like domain family, member 1	0.0071	-2.80
Ssc.8027.1.A1_at	98	dickkopf homolog 3 (Xenopus laevis)	0.0071	5.16
Ssc.5895.1.A1_at	99	collagen, type VI, alpha 1	0.0071	-6.72
Ssc.10793.1.A1_at	100	syndecan 2	0.0071	-10.35
Ssc.20706.1.S1_at	102	hypothetical protein LOC100523635	0.0071	2.94
Ssc.4048.1.A1_at	107	neogenin-like	0.0074	2.43
Ssc.26172.1.S1_at	108	leprecan-like 4	0.0074	-1.01
Ssc.4381.1.S1_at	111	septin 11	0.0077	-8.88
Ssc.820.1.S1_at	113	alanyl (membrane) aminopeptidase	0.0079	-11.29
Ssc.1986.1.S1_at	114	Prostaglandin-endoperoxide synthase 1	0.0079	2.31
Ssc.2425.1.S1_at	115	zonula occludens 1	0.0079	5.21

Table 6. *Differentially expressed genes by depot ($Q < 0.01$) (continued).*

Ssc.11185.1.A1_at	117	SH3 domain-containing kinase-binding protein 1-like	0.0079	-1.01
Ssc.15233.1.S1_at	119	Similar to Homeobox protein SIX1 (Sine oculis homeobox homolog 1)	0.0079	-11.80
Ssc.14134.1.S1_at	120	polypeptide N-acetylgalactosaminyltransferase 1-like	0.0079	-2.70
Ssc.28359.1.A1_at	121	hypothetical protein LOC100153390	0.0079	-0.72
Ssc.7769.2.S1_at	122	lysophosphatidylglycerol acyltransferase 1	0.0080	-7.40
Ssc.1495.1.S1_a_at	125	transcription elongation factor A protein 1-like	0.0081	2.60
Ssc.12446.1.A1_at	127	caspase-13-like	0.0081	-2.16
Ssc.21869.1.S1_at	128	src kinase-associated phosphoprotein 2-like	0.0081	3.60
Ssc.1091.2.S1_at	130	collagen, type I, alpha 1	0.0081	-5.53
Ssc.5895.1.A2_at	132	Collagen, type VI, alpha 1	0.0081	-12.83
Ssc.25217.1.S1_a_at	133	selenoprotein M	0.0081	-6.89
Ssc.11628.1.A1_at	135	GPN-loop GTPase 1-like	0.0081	-1.14
Ssc.15646.1.S1_at	136	lipase maturation factor 2-like	0.0081	-1.66
Ssc.373.1.S1_at	137	matrix Gla protein	0.0081	35.16
Ssc.8072.2.A1_at	139	latent-transforming growth factor beta-binding protein 1-like	0.0081	3.49
Ssc.18454.1.A1_at	143	transducin-like enhancer of split 1 (E(sp1) homolog, Drosophila)	0.0081	2.32
Ssc.3919.1.S1_at	144	cytosolic 5'-nucleotidase III-like protein-like	0.0081	-3.73
Ssc.19566.1.S1_at	147	secreted frizzled-related protein 4	0.0081	-28.97
Ssc.27415.1.S1_at	148	homeobox protein BarH-like 1-like	0.0082	2.65
Ssc.12068.1.A1_at	149	Collagen, type VI, alpha 3	0.0082	-2.95
Ssc.19119.1.A1_at	151	homeobox protein EMX2-like	0.0082	-1.00
Ssc.5645.1.A1_at	152	lamina-associated polypeptide 2, isoforms beta/gamma-like	0.0084	1.58
Ssc.3401.1.S1_at	153	eukaryotic translation initiation factor 3, subunit M	0.0084	2.94
Ssc.11302.1.S2_at	155	hypothetical protein LOC100152001	0.0084	-67.71

Table 6. *Differentially expressed genes by depot ($Q < 0.01$) (continued).*

Ssc.4563.1.S1_a_at	158	N-acetyltransferase 10 (GCN5-related)	0.0086	-0.34
Ssc.6105.1.A1_at	161	plexin B1	0.0087	0.90
Ssc.8000.1.A1_at	166	O-linked N-acetylglucosamine (GlcNAc) transferase (UDP-N-acetylglucosamine:polypeptide-N-acetylglucosaminyl transferase)	0.0091	2.69
Ssc.6736.1.S1_at	168	homeobox protein MSX-2-like	0.0094	-2.07
Ssc.24393.1.S1_at	170	reticulocalbin-3-like	0.0095	-11.50
Ssc.9061.1.A1_at	171	cystatin-M-like	0.0096	2.41
Ssc.3832.1.S1_at	172	aquaporin 3 (Gill blood group)	0.0096	-40.64
Ssc.17281.1.A1_at	174	F-box and leucine-rich repeat protein 5	0.0096	-6.54
Ssc.25178.1.S1_a_at	175	protein transport protein Sec24D-like	0.0096	-2.48
Ssc.8564.1.A1_at	178	Similar to very low density lipoprotein receptor VLDL-R2	0.0096	7.06
Ssc.22738.1.A1_at	179	protein MON2 homolog	0.0096	-1.49
Ssc.2589.1.S1_at	180	troponin I type 3 (cardiac)	0.0096	2.36
Ssc.4993.1.A1_at	181	hypothetical protein LOC100517021	0.0096	-19.82
Ssc.272.1.S1_a_at	182	CD55 molecule, decay accelerating factor for complement (Cromer blood group)	0.0096	1.96
Ssc.26447.1.S1_at	183	carboxypeptidase D	0.0099	1.09
Ssc.18928.1.A1_at	185	Mucosal vascular addressin cell adhesion molecule 1	0.0099	1.37

^aRanking of all differentially expressed ($Q < 0.01$) genes.

^bFold change in mRNA abundance for visceral adipose tissue relative to subcutaneous adipose tissue. A positive fold change indicates increased mRNA abundance in visceral adipose tissue relative to subcutaneous adipose tissue and a negative fold change indicates decreased mRNA abundance.

^cResiduals of gene showed evidence of a non-normal distribution

Table 7. *Gene ontology cellular component: extracellular region gene list^a.*

Affy ID	Gene Title	Q-value	Fold Change
Ssc.12365.1.A1_at	ADAM metallopeptidase with thrombospondin type 1 motif, 1	0.0233	-6.48
Ssc.11257.1.S1_at	TIMP metallopeptidase inhibitor 2	0.0130	-34.61
Ssc.6009.1.S1_at	amylase, alpha 2B (pancreatic)	0.0440	-0.58
Ssc.17345.1.S1_at	angiopoietin-like 4	0.0068	-2.56
Ssc.14474.1.S1_at	Apomucin	0.0206	0.09
Ssc.16624.1.S1_at	carbonic anhydrase-related XI protein	0.0342	0.58
Ssc.1020.1.S1_at	cathepsin K	0.0170	-28.91
Ssc.657.1.A1_at	chemokine (C-C motif) ligand 2	0.0327	-4.05
Ssc.18613.1.S1_at	chemokine (C-C motif) ligand 21	0.0180	1.94
Ssc.4871.1.S1_at	chemokine (C-X-C motif) ligand 2	0.0215	-2.08
Ssc.31199.1.S1_at	collagen, type V, alpha 1	0.0179	-0.80
Ssc.61.1.S1_at	complement component 3	0.0345	31.49
Ssc.10245.2.A1_a_at	Decorin	0.0281	-41.25
Ssc.8027.1.A1_at	dickkopf homolog 3	0.0071	5.16
Ssc.16236.1.S1_at	dipeptidyl-peptidase 4	0.0489	0.67
Ssc.15923.1.S1_at	fibroblast growth factor 7 (keratinocyte growth factor)	0.0039	-0.25
Ssc.137.1.S1_at	Ficolin	0.0184	-14.97
Ssc.5833.1.A2_at	follicle stimulating hormone, beta polypeptide	0.0222	0.05
Ssc.4747.1.S1_at	Follistatin	0.0168	0.63
Ssc.5713.1.S1_at	gelatinase A	0.0015	-50.51
Ssc.14246.1.S1_at	Gelsolin	0.0302	-23.34
Ssc.16231.1.S1_a_at	insulin-like growth factor 1	0.0119	-2.68
Ssc.15588.1.S2_at	insulin-like growth factor binding protein 3	0.0339	-16.60
Ssc.15800.1.S1_at	insulin-like growth factor binding protein 5	0.0302	-10.14
Ssc.14062.2.S1_a_at	insulin-like growth factor binding protein 6	0.0281	-20.02
Ssc.62.2.S1_a_at	interleukin 6 (interferon, beta 2)	0.0470	-0.47
Ssc.373.1.S1_at	matrix Gla protein	0.0081	35.16
Ssc.734.1.S1_at	matrix metallopeptidase 14 (membrane-inserted)	0.0418	-2.07
Ssc.11194.1.S1_at	plasminogen activator	0.0383	-2.24
Ssc.196.1.S1_at	plasminogen activator, tissue	0.0296	-6.52

Table 7. *Gene ontology cellular component: extracellular region gene list^a(Continued).*

Ssc.16228.1.S1_at	platelet basic protein	0.0417	1.73
Ssc.10510.1.A1_at	Prolactin	0.0480	-0.12
Ssc.14556.1.S1_at	Proopiomelanocortin	0.0136	0.10
Ssc.15695.1.S1_at	retinol binding protein 4, plasma	0.0206	19.02
Ssc.2464.1.S1_at	stanniocalcin 1	0.0040	-4.30
Ssc.5614.1.S1_at	Tryptase	0.0413	0.49

^aGene list of over-represented family of genes (EASE Score=0.0050) that were differentially expressed (Q<0.05).

^bFold change in mRNA abundance for visceral adipose tissue relative to subcutaneous adipose tissue. A positive fold change indicates increased mRNA abundance in visceral adipose tissue relative to subcutaneous adipose tissue and a negative fold change indicates decreased mRNA abundance.

Table 8. KEGG pathway: arachidonic acid metabolism gene list^a.

Affy ID	Gene Title	Q-value	Fold Change
Ssc.10974.1.S1_at	arachidonate 12/15-lipoxygenase	0.0124	13.64
Ssc.732.1.S1_at	carbonyl reductase 2	0.0461	3.06
Ssc.11267.1.S1_at	cytochrome P450 2B22	0.0424	2.95
Ssc.1310.1.S1_at	microsomal prostaglandin E synthase-1	0.0070	-4.09
Ssc.7314.1.A1_at	prostaglandin G/H synthase-2	0.0195	-0.12
Ssc.1986.1.S1_at	prostaglandin-endoperoxide synthase 1	0.0079	2.31
Ssc.8278.1.S1_at	soluble epoxide hydrolase	0.0264	-4.27

^aGene list of over-represented family of genes (EASE Score=0.0152) that were differentially expressed (Q<0.05).

^bFold change in mRNA abundance for visceral adipose tissue relative to subcutaneous adipose tissue. A positive fold change indicates increased mRNA abundance in visceral adipose tissue relative to subcutaneous adipose tissue and a negative fold change indicates decreased mRNA abundance.

Table 9. KEGG pathway: chemokine signaling pathway gene list^a.

Affy ID	Gene Title	Q-value	Fold Change ^b
Ssc.657.1.A1_at	chemokine (C-C motif) ligand 2	0.0327	-4.04524
Ssc.18613.1.S1_at	chemokine (C-C motif) ligand 21	0.018	1.938086
Ssc.19054.1.S1_a_at	chemokine (C-C motif) receptor 3	0.0488	0.169559
Ssc.4871.1.S1_at	chemokine (C-X-C motif) ligand 2	0.0215	-2.07684
Ssc.14176.1.A1_s_at	glycogen synthase kinase 3 beta	0.015	-1.22895
Ssc.16228.1.S1_at	platelet basic protein	0.0417	1.727623
Ssc.6025.1.S1_at	signal transducer and activator of transcription 1	0.0311	1.367214
Ssc.10911.2.S1_at	similar to Beta-adrenergic receptor kinase 2 (Beta-ARK-2) (G-protein-coupled receptor kinase 3)	0.0143	-3.14621
Ssc.2140.1.S1_at	similar to PTK2B protein tyrosine kinase 2 beta	0.0471	0.422968
Ssc.5093.1.A1_at	similar to SHC-transforming protein 1 (SH2 domain protein C1) (Src homology 2 domain-containing-transforming protein C1)	0.0407	-3.14165

^aGene list of over-represented family of genes (EASE Score=0.0152) that were differentially expressed (Q<0.05).

^bFold change in mRNA abundance for visceral adipose tissue relative to subcutaneous adipose tissue. A positive fold change indicates increased mRNA abundance in visceral adipose tissue relative to subcutaneous adipose tissue and a negative fold change indicates decreased mRNA abundance.

CHAPTER 6. GENERAL CONCLUSIONS

The prevalence of obesity in the United States is continuing to rise (2-3) and places a significant burden on our health care system (11). Research has identified that insulin resistance in obesity is related to adipose tissue inflammation (4) and that this is accompanied by macrophage infiltration into adipose tissue (12-13). Rodent models have been pivotal to obesity and obesity-related research and the role of individual genes. However, distinct differences between rodents and humans underscore the need for alternative comparative models. Unlike humans, CRP is not an acute phase protein nor is it highly expressed in mice (10). Additionally, mice macrophages are functionally different in that they produce NO via inducible nitric oxide synthase (8). Swine are an attractive alternative model due to their comparative physiology and anatomy to humans. In particular, Ossabaw swine are a suitable alternative as they can consistently develop multiple metabolic syndrome risk factors (9).

I first sought to characterize obesity-linked adipose tissue inflammation and macrophage phenotype in Ossabaw swine (Chapter 3). Therefore, I feed Ossabaw swine for 30 weeks with a restricted intake low-fat control diet, a high palm oil diet, or a high palm oil with n-3 fatty acids diet. We determined that the major adipose tissue macrophage phenotype in Ossabaw swine is $CD16^+CD14^-$, which is likely anti-inflammatory. High dietary palm oil did cause an increase in adipose tissue $CD16^-CD14^+$ macrophages, which are likely inflammatory, and this increase was attenuated by the addition of dietary n-3 fatty acids. Concomitant with a largely anti-inflammatory adipose tissue macrophage phenotype, there was no induction of inflammation measured by serum CRP, $TNF\alpha$, IL-6, or IL-12. Serum adiponectin was decreased by the high palm oil diet, but this decrease was attenuated by the addition of n-3 fatty acids. However, the increase in serum adiponectin by the

addition of n-3 fatty acids did not attenuate the decrease in longissimus dorsi AMPK α phosphorylation. Some individuals remain healthy despite obesity, and possess a favorable inflammation profile (5). The ability to expand adipose tissue may afford protection against adipose tissue expansion. As evidenced by Kim *et al.* (6), adipose tissue expansion attenuates inflammation due to obesity.

However, the metabolic profile of obese Ossabaw swine in this study was not entirely healthy. There was no increase in serum cholesterol or triglycerides, but there was apparent hyperglycemia without hyperinsulinemia. The lack of hyperinsulinemia suggests that there was impaired β -cell function in the obese Ossabaw swine, which could be caused by a number of obesity related complications including ectopic lipid deposition in the pancreas. Although there was no ectopic lipid deposition in the liver or longissimus dorsi muscle. Challenging Ossabaw swine with insulin revealed that high dietary palm oil induced insulin resistance, but the addition of dietary n-3 fatty acids attenuated the palm oil-induced insulin resistance. Despite the hyperglycemia and insulin resistance, the capacity to markedly expand adipose tissue may have afforded protection against obesity-linked inflammation. This demonstrates, at least in this study, that inflammation is not definitely linked to inflammation, at least in the parameters we measured.

Due to inflammation being largely absent, we hypothesized that the ability to expand adipose tissue protected Ossabaw swine against adipose tissue inflammation. Therefore, we determined whether SVC-derived adipocytes from Ossabaw swine were protected from palmitate-induced inflammation and insulin resistance compared to Yorkshire SVC-derived adipocytes (Chapter 4). We determined that NF κ B was equally activated in both Ossabaw and Yorkshire adipocytes in response to palmitate. However, only Yorkshire adipocytes

secrete the chemokine IL-8 in response to palmitate treatment. IL-8 is responsible for recruiting neutrophils to the site of injury and inflammation, and are present in adipose tissue after one week of high fat feeding in mice (1). The absence of palmitate-induced IL-8 production by Ossabaw adipocytes may be involved in the lack of substantial adipose tissue inflammation seen in our first study. Consistent with the lack of palmitate-induced IL-8 production in adipocytes, there was no increase in serum or adipose tissue IL-8 in obese Ossabaw swine.

Palmitate decreased insulin-stimulated glucose uptake in both adipocyte genotypes. However, Ossabaw adipocytes had decreased basal and insulin-stimulated glucose uptake compared to Yorkshire adipocytes. Although the fold increase for insulin-stimulated glucose uptake was equal between the two adipocyte genotypes. Though there have been no direct comparisons, this *in vitro* data matches two separate *in vivo* insulin challenges in Ossabaw and Yorkshire swine, where Yorkshire swine comparably have more insulin sensitivity than Ossabaw swine. Whether this decreases overall insulin sensitivity of Ossabaw swine contributes to the susceptibility of Ossabaw swine to develop insulin resistance remains to be determined.

Because of the potential for adipose expansion to protect against inflammation, we performed microarray analysis on subcutaneous and visceral adipose tissue of lean and obese Ossabaw swine (Chapter 5). Therefore, Ossabaw swine were fed either a restricted or an *ad libitum* nutritionally adequate diet for 38 weeks. Microarray analysis revealed no dietary treatment induced differences in gene expression after controlling for FDR. Coinciding with this, there were no negative obesity-linked metabolic alterations. The lack of a high fat diet is likely causative to the lack of metabolic alteration in obese Ossabaw swine. The

importance of high fat in inducing metabolic alterations is evident in the fructose group of Lee *et al.* (7) which did not cause a negative metabolic profile. There were 1494 differently expression genes by depot after controlling for FDR, of which 777 were annotated. One gene ontology cellular component and two KEGG pathways were determined to contain over-represented genes from our significantly expressed genes by depot. These were extracellular region, arachidonic acid metabolism, and chemokine signaling pathway, respectively.

Overall, we have demonstrated that Ossabaw swine have a favorable inflammatory profile in response to a high fat diet, despite morbid obesity. Additionally, the inflammatory profile was not linked to negative metabolic consequences. Potentially, the ability to expand adipose tissue protects against inflammation, which is supported by palmitate induced IL-8 induction in Yorkshire, but not Ossabaw adipocytes. This is further evidenced by the lack of obesity-linked microarray gene expression differences in Ossabaw swine when fed a diet without high fat.

This research can be furthered in multiple areas. Perhaps most intriguing is whether the decreased basal and insulin-stimulated glucose uptake in Ossabaw adipocytes versus Yorkshire adipocytes is a component of the ability to Ossabaw swine to develop insulin resistance. Additionally, genetic determination of the cause of the reduced glucose uptake and whether analogous conditions is a component of obesity-linked insulin resistance in humans. Additionally, continued research into the capacity of Ossabaw swine to expand adipose tissue and its ability to afford protection against obesity-linked inflammation is an intriguing question, particularly if it is analogous to the metabolically healthy obese human population subset. Finally, because of immunological differences between humans and mice,

further characterization of swine adipose tissue macrophages with regards to phenotype and function is an imminent research question to further the validation of the model for obesity and obesity-linked research.

References

1. Elgazar-Carmon V, Rudich A, Hadad N, Levy R. 2008. Neutrophils transiently infiltrate intra-abdominal fat early in the course of high-fat feeding. *J Lipid Res* 49:1894-1903.
2. Flegal KM, Carroll MD, Ogden CL, Curtin LR. 2010. Prevalence and trends in obesity among US adults, 1999-2008. *JAMA* 303:235-241.
3. Flegal KM, Carroll MD, Ogden CL, Johnson CL. 2002. Prevalence and trends in obesity among US adults, 1999-2000. *JAMA* 288:1723-1727.
4. Hotamisligil GS, Shargill NS, Spiegelman BM. 1993. Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance. *Science* 259:87-91.
5. Karelis AD, Faraj M, Bastard JP, St-Pierre DH, Brochu M, Prud'homme D, Rabasa-Lhoret R. 2005. The metabolically healthy but obese individual presents a favorable inflammation profile. *J Clin Endocrinol Metab* 90:4145-4150.
6. Kim JY, van de Wall E, Laplante M, Azzara A, Trujillo ME, Hofmann SM, Schraw T, Durand JL, Li H, Li G, Jelicks LA, Mehler MF, Hui DY, Deshaies Y, Shulman GI, Schwartz GJ, Scherer PE. 2007. Obesity-associated improvements in metabolic profile through expansion of adipose tissue. *J Clin Invest* 117:2621-2637.
7. Lee L, Alloosh M, Saxena R, Van Alstine W, Watkins BA, Klaunig JE, Sturek M, Chalasani N. 2009. Nutritional model of steatohepatitis and metabolic syndrome in the Ossabaw miniature swine. *Hepatology* 50:56-67.
8. Mestas J, Hughes CC. 2004. Of mice and not men: differences between mouse and human immunology. *J Immunol* 172:2731-2738.
9. Neeb ZP, Edwards JM, Alloosh M, Long X, Mokelke EA, Sturek M. 2010. Metabolic syndrome and coronary artery disease in Ossabaw compared with Yucatan swine. *Comp Med* 60:300-315.
10. Szalai AJ, McCrory MA. 2002. Varied biologic functions of C-reactive protein: lessons learned from transgenic mice. *Immunol Res* 26:279-287.

11. Wang YC, McPherson K, Marsh T, Gortmaker SL, Brown M. 2011. Health and economic burden of the projected obesity trends in the USA and the UK. *Lancet* 378:815-825.
12. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW, Jr. 2003. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* 112:1796-1808.
13. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS, Tartaglia LA, Chen H. 2003. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest* 112:1821-1830.